

# The Role of Friends in the Opioid Epidemic<sup>\*†</sup>

Effrosyni Adamopoulou<sup>1</sup>, Jeremy Greenwood<sup>2</sup>, Nezih Guner<sup>3</sup>, and Karen Kopecky<sup>4</sup>

<sup>1</sup>ZEW, University of Mannheim, and IZA

<sup>2</sup>University of Pennsylvania

<sup>3</sup>CEMFI and Banco de España

<sup>4</sup>FRB Cleveland and Emory University

August 25, 2025

## Abstract

The role of friends in the US opioid epidemic is examined. Using data from the National Longitudinal Survey of Adolescent Health (Add Health), adults aged 25-34 and their high school best friends are focused on. An instrumental variable technique is employed to estimate peer effects in opioid misuse. Severe injuries in the previous year are used as an instrument for opioid misuse in order to estimate the causal impact of a person's best friends' opioid misuse on their own misuse. The estimated peer effects are significant: Having a best friend who misuses opioids following a serious injury increases the probability of own opioid misuse by around 7 percentage points in a population where 17 percent ever misuses opioids. The effect is concentrated among non-college graduates and peers with strong ties who are central in their friendship networks. Peer opioid misuse eventually leads to deteriorating health and opioid addiction.

JEL: C26, D10, I12, J11.

Keywords: opioid, peer-group effects, friends, instrumental variables, Add Health, severe injuries.

<sup>\*</sup>We thank Jiangting Wang for excellent research assistance and Manuel Arellano, Dmitry Arkhangelsky, Michèle Belot, Christopher Cronin, Bill Dowd, Joseph Sabia, Hannes Schwandt, Bryce J. Stanley, Konstantinos Tatsiramos, Jenny Williams, the participants at the public economics seminar at the University of Mannheim, the DEM Research Seminar at the University of Luxembourg, the EuHEA Online Seminar, the 2024 Essen Health Conference, the 4th Workshop on Networks & Development in Naples, the 2024 Economics of Mental Health Workshop in Minneapolis, the seminar at VCU, the Collaborative Health Economics Seminar Series at GW, and at the UC3M alumni conference for useful comments and suggestions. This research uses data from Add Health, funded by grant P01 HD31921 (Harris) from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), with cooperative funding from 23 other federal agencies and foundations. Add Health is currently directed by Robert A. Hummer and funded by the National Institute on Aging cooperative agreements U01 AG071448 (Hummer) and U01AG071450 (Aiello and Hummer) at the University of North Carolina at Chapel Hill. Add Health was designed by J. Richard Udry, Peter S. Bearman, and Kathleen Mullan Harris at the University of North Carolina at Chapel Hill. Information on how to obtain the Add Health data files is available on the Add Health website (<https://addhealth.cpc.unc.edu>). No direct support was received from grant P01-HD31921 for this project. Adamopoulou acknowledges financial support from the German Research Foundation through CRC TR 224 (project A03). The views expressed herein are those of the authors and not necessarily those of Banco de España, the Eurosystem, the Federal Reserve Bank of Cleveland, or the Federal Reserve System.

<sup>†</sup>Contact info: Effrosyni Adamopoulou, [effrosyni.adamopoulou@zew.de](mailto:effrosyni.adamopoulou@zew.de)

# 1 Introduction

Opioids have led to the worst drug overdose epidemic in US history—see [Cutler and Glaeser \[2021\]](#) and [Maclean et al. \[2022\]](#) for recent reviews—and have had a deleterious impact on public health and individual labor-market outcomes. What role have peers played in the spread of the opioid crisis? For those who misuse prescription opioids, friends are one of the most common sources for obtaining them. Table 1, panel A, shows the fraction of individuals between the ages of 26 and 34 who misuse opioids and get them from friends and relatives. Between 2010 and 2019, more than 50 percent of opioid misusers obtained opioids from friends or relatives. In 2010, when the total opioid prescription shipments peaked in the United States, the role of friends and relatives was even stronger, possibly due to the wider availability of prescribed opioids. This widespread reliance on social networks for opioid acquisition implies that peer effects are likely to be substantial drivers in the diffusion of opioid misuse. Hence, it is not surprising that peers have been highlighted as a potentially important factor in the fight against opioids [[Compton et al., 2019](#), [Blanco et al., 2020](#)]. Yet, the empirical estimation of peer effects is challenging due to the unavailability of data and the difficulties in achieving identification.

Table 1: Individuals ages 26-34 who misuse prescription opioids:  
main method of acquiring and main reason for misuse

| Year  | All    | Non-College | College |
|---|--------|-------------|---------|
| <i>Panel A: Fraction that obtained opioids from friends</i>             |        |             |         |
| 2010  | 56.35% | 56.16%      | 56.91%  |
| 2015  | 46.32% | 46.15%      | 46.91%  |
| 2019  | 34.83% | 33.26%      | 38.95%  |
| <i>Panel B: Physical pain as main reason for last misuse of opioids</i> |        |             |         |
| 2015  | 59.14% | 61.52%      | 58.42%  |

**Note:** Calculations based on data from the National Survey on Drug Use and Health (NSDUH), an annual nationwide survey that provides national and state-level data on the use of tobacco, alcohol, illicit drugs (including the non-medical use of prescription drugs), and mental health in the United States. The misuse of prescription drugs is defined as use in any way that is not directed by a doctor during the last 12 months—i.e., without a prescription, use in greater amounts than prescribed, more often than prescribed, longer than prescribed, or in any other non-directed way.

This gap is filled here by providing causal evidence on peer effects in opioid misuse using a novel identification strategy that exploits quasi-random exposure to opioid prescriptions following severe injuries among best friends. The analysis draws on rich information on opioid misuse among individuals aged 25–34 and their high school best friends, leveraging comprehensive controls for demographics, health, and parental characteristics. A range of exercises, sample exclusions, and placebo tests are implemented to address concerns

related to selection, homophily, and potential violations of the exclusion restriction.

The analysis uses the National Longitudinal Study of Adolescent to Adult Health (Add Health), a longitudinal study of a nationally representative sample of over 20,000 adolescents who were in grades 7-12 during the 1994-95 school year, and have been followed for five waves to date, most recently in 2016-18. The main analysis draws data from Wave IV of Add Health in 2009; i.e., at the first stage of the opioid epidemic, when knowledge regarding the risk of addiction and death due to opioid misuse was still limited and the prescriptions for opioids as painkillers were very common—see [Guy \[2017\]](#) and [Figure A1](#).<sup>1</sup> For example, [Muench et al. \[2020\]](#) report that opioid prescriptions in U.S. community health centers serving low-income patients decreased by 73.7% from 2009 to 2018, highlighting the high prevalence of prescribing in 2009 and its decline over time. While first heroin and then synthetic opioids (mainly fentanyl) played an increasingly significant role in the opioid crisis in more recent years (see [Figure A1](#)), the period studied here primarily reflects misuse of prescribed opioids, which were the dominant driver of opioid misuse at that time. In the analysis, opioid misuse refers to using prescription painkillers like Vicodin or OxyContin without a prescription, in larger amounts, more often, longer than prescribed, or for the experience they cause.

The starting point for the analysis is the evidence that severe injuries sustained by best friends have a significant effect on an individual’s opioid misuse. In an ordinary least squares (OLS) regression, having a best friend who experiences a severe injury increases an individual’s own probability of opioid misuse by approximately 7.3 percentage points (pp). This effect is substantial, about 80 percent of the estimated effect of experiencing a severe injury oneself, suggesting a large role for peer diffusion in opioid misuse.

What explains this link between a friend’s injury and an individual’s opioid misuse? The proposed mechanism is peer effects. Individuals can develop opioid misuse or addiction after being prescribed opioids for a severe injury—an exogenous event—which can then lead to misuse among their peers. Prior research shows that even a single exposure to opioids, such as during an emergency room visit or a C-section, can significantly increase the risk of future misuse, even months later [[Barnett et al., 2017](#), [Eichmeyer and Zhang, 2022](#), [Carrico et al., 2020](#)].<sup>2</sup> Supporting this, as shown in [Table 1](#), panel B, 59 percent of opioid misusers

<sup>1</sup>Deaths due to opioid misuse increased sharply and steadily after 2014. The last part of this paper analyzes the welfare implications of peer opioid misuse on the probability of addiction and death.

<sup>2</sup>[Currie and Zwiens \[2023\]](#) similarly find that women prescribed antidepressants postpartum often continue using them long-term, highlighting how an initial medical prescription—though exogenous—can lead to sustained medication use.

aged 26–34 cite physical pain as the primary reason for their misuse.

To identify the causal effect of peer influence, the analysis employs an instrumental variable (IV) approach, controlling for state and/or school-specific factors. Using best friends’ severe injuries as an instrument for their opioid misuse reveals a strong positive peer effect. Having a best friend who misuses opioids after a serious injury raises the probability of own opioid misuse by around 7 pp in a population where 17 percent have misused opioids. The estimated effect is close to that found in the reduced form analysis.

Several potential threats to identification are addressed. First, the reduced-form evidence on effects of friend’s injury on opioid misuse can be due to a general tendency toward risky behavior among friends or correlated effects from shared environments, such as local conditions or policies affecting both individuals and their peers. However, in a placebo analysis, there is no effect of best friends’ injuries on the probability of smoking, getting drunk, or having unprotected sex, other forms of risky behavior. Severe injuries by non-best friends who live in the same county also do not generate any effect on opioid misuses. Moreover, controlling for a wide range of factors, including parental drug availability, predetermined friend characteristics, Big Five personality traits, risk aversion, occupational status, and socioeconomic background, leaves the IV results unchanged, and further elevating concerns for potential correlated effects.

Second, the validity of the IV strategy hinges on the assumption that friends’ injuries are exogenous, affecting opioid misuse only through the friend’s own misuse. Yet, injuries may directly cause injuries in others (e.g., via shared accidents). Opioid misuse might also be due to stress caused by the injuries of best friends directly, and not by their opioid use, again potentially violating the exclusion restriction. The results remain robust, however, when the analysis is restricted to injuries unrelated to vehicle accidents, reducing the risk of shared injury events. They are also robust when controls are added for whether an individual was diagnosed by depression, post-traumatic stress disorder or anxiety, potential factors that can lead to opioid misuse. Additionally, there is no significant correlation between own injuries and friends’ injuries, supporting the validity of the exclusion restriction.

Finally, the IV analysis assumes that, once we control for observables, the instrument (the severe injuries) is random with respect to other unobserved determinants of opioid misuse. Several sample restrictions are applied to minimize the potential confounding effects of unobserved traits. In particular, the results hold

when the analysis is restricted to individuals and their best friends who have no prior experience with illegal drugs or non-prescription painkillers, or to those who are not identified as risk-takers. They also hold when individuals with severe injuries themselves are excluded, as their own injuries could directly lead to opioid misuse through prescribed opioids.

In the main analysis, a person is considered to misuse opioids if they report that they ever misuse opioids. This can raise concerns about the timing, as opioid misuse might not necessarily follow the injuries of best friends. Also, misuse is a simple yes/no variable, without an indication of intensity of misuse. To address these concerns, the analysis also uses measures of opioid misuse frequency over the past 12 months. The results remain robust across different levels of misuse intensity, from occasional use to more frequent consumption (e.g., 1–2 days per week), highlighting the persistence of peer influence on both the initiation and escalation of opioid misuse.

The results show that the peer effect is concentrated among individuals without a college degree, and among close, long-lasting friendships, particularly same-gender friends residing in the same county. The effect is amplified in dense or large friendship networks and among individuals who occupy central positions within their social networks. Peer effects are also stronger in areas with high opioid dispensing rates, delayed implementation of naloxone access laws, higher prevalence of physical and mental health issues, limited healthcare access, and low social capital.

Finally, the analysis reveals significant adverse health and welfare consequences of peer opioid misuse. Individuals exposed to peer misuse report worse self-assessed health, a higher likelihood of stimulant misuse, and a greater probability of opioid addiction. There is also suggestive, but imprecisely estimated, evidence of a link between peer opioid misuse and increased risk of death from drug poisoning or suicide.

**Related Literature.** This study contributes to the recent literature on the determinants of the opioid epidemic in the United States (see, among others, [Alpert et al. 2018](#); [Ruhm 2019](#); [Alpert et al. 2022](#); [Eichmeyer and Zhang 2022](#); [Finkelstein et al. 2022](#); [Dowd 2023](#); [Eichmeyer and Zhang 2023](#); [Janssen and Zhang 2023](#)).<sup>3</sup> The analysis focuses on the first stage of the epidemic and appears to be the first paper that causally identifies, at the micro level, the role of friends in its spread in the United States. Previous studies, such

<sup>3</sup>[Finkelstein et al. \[2022\]](#) find an important role for location-specific factors, which can potentially capture differences in peer effects across locations.

as [Mäckle and Ruenzi \[2022\]](#) and [Cutler and Donahoe \[2024\]](#), have examined network effects on overdose deaths using friendship links—such as Facebook connections—at the county level and have employed panel data to study the evolution of opioid deaths over time. In contrast, the current investigation offers a more granular analysis by focusing on opioid misuse and addiction at the individual level during the initial phase of the epidemic. It looks at long-established friendships formed during adolescence, intentionally excluding newly formed friendships that may be more prone to endogeneity concerns. The large effect found in this study aligns with these studies at the county level, which show that spillovers between geographically close counties or those linked through Facebook networks can account for 84 to 92 percent of opioid deaths from 1990 to 2018, highlighting the significant role of network effects in the opioid epidemic. The current study’s individual-level approach provides a more detailed understanding of how peer relationships specifically contribute to the spread of opioid misuse and addiction.

[Seamans et al. \[2018\]](#) study opioid initiation among household members using a similar instrument (injury of a family member), distinguishing by injury type (ankle sprain or fracture, with the latter being treated more frequently with opioids). The estimated positive effect on other household members highlights the relevance of drug availability at home but is difficult to interpret as a peer effect due to homophily and correlated effects among family members.<sup>4</sup> The strategy here is to estimate peer effects among high school best friends including state and/or school fixed effects to account for correlated effects. [Cappellari and Tatsiramos \[2015\]](#) use the onset of best friends’ health problems as an instrument of their employment status to examine peer effects on job finding. The proposed instrument here, that is, best friends’ severe injuries, is conceptually similar but is used to study peer effects on opioid misuse. Finally, [Thingholm \[2023\]](#) documents spillovers in opioid prescriptions among practitioners in Denmark and the negative consequences on their patients’ labor market outcomes, while [Rose et al. \[2024\]](#) examine the influence of peers on recovery from substance use disorders among patients in Norway.

The findings are also related to the literature that studies peer effects on the consumption of other substances such as tobacco or alcohol [[Card and Giuliano, 2013](#), [Cutler and Glaeser, 2005](#), [Clark and Lohéac, 2007](#), [Eisenberg et al., 2014](#), [Kremer and Levy, 2008](#), [Lundborg, 2006](#), [Fletcher, 2012](#)]. Some of these studies

<sup>4</sup>See also [[de Vaan and Stuart, 2019](#), [Khan et al., 2019](#)] for a similar approach, but with similar challenges in interpretation.

also adopt an instrumental variable technique, using substance availability at friends’ parental homes or average characteristics of friends as instruments. The current analysis contributes by focusing on opioid misuse 14 years after friendship formation (not contemporaneous) and by proposing an instrument whose exclusion restriction is more likely to hold. Beyond substance abuse, the instrumental variables approach is widespread in the literature that causally estimates peer effects—see among others [Dahl et al. \[2014\]](#), [De Giorgi et al. \[2010\]](#), [Joensen and Nielsen \[2018\]](#), and [Kim et al. \[2024\]](#).

## 2 Data

To analyze peer effects on opioid misuse, unique information is garnered from Add Health on best friends in high school and subsequent opioid misuse in adulthood (14 years after). Add Health is a longitudinal survey of a nationally representative sample of over 20,000 adolescents who were in grades 7-12 during the 1994-95 school year—see [Figure A2](#) and [Harris \[2018\]](#). Information is harnessed from four waves. Longitudinal survey weights are used to account for any possible attrition. More specifically, Wave I of the survey took place in 1994/1995 and entailed in-home interviews of a representative sample of high school students in the United States. Respondents were asked to nominate up to five male and five female friends. Nominations were made starting from the closest friend to the most distant friend. The focus is on the first male and female nominations; i.e., the best friends. This choice is motivated by the higher likelihood of the respondents staying in contact with best friends in adulthood and the low fraction of respondents, less than 1/3 of them, nominating more than two friends. Respondents were asked to nominate their friends again in Wave II, about a year after Wave I. This information is used to analyze heterogeneity in the effects among friendships of different duration. Given that, in most cases, individuals and their best friends were attending the same school, they were all part of the Add Health in-home interview, which allows the retrieval of a rich set of information for both the individuals and their best friends.<sup>5</sup>

Friends nominated in the Add Health study can be used to map out friendship networks within each high school, and Add Health also provides detailed information about these networks and each person’s role within

<sup>5</sup>Friendship networks are used to study peer effects on different socioeconomic outcomes, such as education, living arrangements, and teenage pregnancies (see, among others, [Bifulco et al. 2011](#); [Fernández-Villaverde et al. 2014](#); [Patacchini et al. 2017](#); [Adamopoulou and Kaya 2018](#); [Agostinelli et al. 2022](#)).

them. These measures include: i) Bonacich centrality, defined as the centrality of an individual weighted by the centrality of those they nominate as friends—see Bonacich [1987]; ii) a “reach” measure, representing the maximum number of connections a node (an individual) can connect within the entire friendship network; and iii) relative density, calculated as the ratio of actual ties to the maximum possible ties, computed under the assumption that each respondent could nominate up to 10 friends. These metrics help explore how different network characteristics affect outcomes.

Besides providing information on friendship formation, Wave I of Add Health also contains information on several socioeconomic, educational, and behavioral outcomes for teenagers and their families. In particular, it contains information on the availability of cigarettes and alcohol in respondents’ homes. There is also information on the availability of drugs at home and on whether the respondent had consumed any illegal drug by Wave I. Last, for a subset of respondents whose parents participated in the parents’ interview, there is information on whether the respondent lived with both parents and questions about maternal education and household gross income.

After Waves I and II, respondents (individuals and their friends) were followed to adulthood. Wave III took place in 2001 and Wave IV took place in 2008. A question that allows the *direct* measurement of opioid misuse was asked for the first time in Wave IV.<sup>6</sup> At that time, the respondents were between 25 and 34 years old. The question was:

“Which of the following types of prescription drugs have you ever taken that were not prescribed for you, taken in larger amounts than prescribed, more often than prescribed, for longer periods than prescribed, or that you took only for the feeling or experience they caused? Painkillers or opioids, such as Vicodin, OxyContin, Percocet, Demerol, Percodan, or Tylenol with codeine.”<sup>7</sup>

Additionally, another question in Wave IV restricts the reference period to the last 12 months and assesses the intensity of the misuse: “During the past 12 months, on how many days did you use painkillers when they were not prescribed for you, in larger amounts than prescribed, more often than prescribed, or for longer periods than prescribed?: none, 1 or 2 days, once a month or less, 2 or 3 days a month, 1 or 2 days a

<sup>6</sup>In Wave I respondents were asked about illegal drug use and in Wave III about the use of painkillers (Darvon, Demerol, Percodan, or Tylenol with codeine) without doctor’s permission. Respondents who reported illegal drug use in Wave I or unsubscribed painkiller use in Wave III are excluded in a battery of robustness checks.

<sup>7</sup>The definition of misuse of prescription drugs is very similar in the NSDUH and Add Health. A slightly different age bracket is used in Table 1 since age is reported in particular brackets in the NSDUH.



week, 3 to 5 days a week, every day or almost every day.”

Respondents in Wave IV were also asked to report their self-perceived health status, as well as stimulants, and other drug use. Additionally, there was a question to assess opioid addiction, which asked: “Have you ever continued to use painkillers after you realized using them was causing you any emotional problems (such as feeling depressed or empty, feeling irritable or aggressive, feeling paranoid or confused, feeling anxious or tense, being jumpy or easily startled) or causing you any health problems (such as heart pounding, headaches or dizziness, or sexual difficulties)?”.

Importantly, respondents in Wave IV were asked whether they had suffered a serious injury in the previous year. This question is key for the construction of the instrument and asked: “In the past 12 months, have you suffered any serious injuries? For example, broken bones, cuts or lacerations, burns, torn muscles, tendons or ligaments, or other injuries that interfered with your ability to perform daily tasks.”

Moreover, the Individual Vital Status and Underlying Cause of Death file provides the vital status of each Add Health sample member from Wave I through 2021 and the underlying International Classification of Diseases causes of death code for each decedent. Given that deaths in Add Health are rare events (respondents were between 37 and 46 years old in 2021), the causes of death are reported in aggregated categories. For the current analysis, deaths are then classified into two broad categories, namely, deaths due to medical factors (e.g., HIV, cancer, diabetes, CVD, parasitic, respiratory, or digestive diseases) and deaths due to poisoning from toxic substances or suicide. The latter include drug overdoses, which are likely to be due to opioid misuse.<sup>8</sup>

As Table 2, column 2, shows, almost 17 percent of individuals reported misusing opioids and 14 percent reported a serious injury during the last year, according to data from Wave IV.<sup>9</sup> The reference period for this question is 2008. This is a period when the opioid dispensing rate in the United States was high (above 75 prescriptions per 100 persons) and increasing.<sup>10</sup> Additionally, 3.5 percent of the respondents reported

<sup>8</sup>The Add Health category that contains deaths due to poisoning from toxic substances also includes deaths due to drowning, a number likely to be very small.

<sup>9</sup>Between 2005 and 2019, among individuals between the ages of 26 and 34, about 7 percent report misusing opioids during the last 12 months in the NSDUH. This percentage is lower than in Add Health as the reference period is “the last 12 months” in the NSDUH questionnaire as opposed to “ever taken” in the Add Health questionnaire.

<sup>10</sup>The opioid dispensing rate per 100 persons is defined as the ratio of the total number of prescriptions dispensed annually at the national level over the annual resident population. In the United States, it peaked in 2012 (reaching 81) and subsequently dropped (down to 43 in 2020). See <https://www.cdc.gov/overdose-prevention/data-research/facts-stats/us-dispensing-rate-maps.html> and <https://www.cdc.gov/mmwr/volumes/66/wr/mm6626a4.htm>.

Table 2: Final sample statistics

|   | N<br>(1) | mean<br>(2) | sd<br>(3) | min<br>(4) | max<br>(5) |
|---|----------|-------------|-----------|------------|------------|
| Outcomes (Wave IV)                                      |          |             |           |            |            |
| Opioid misuse   | 2,826    | 0.169       | 0.375     | 0          | 1          |
| Any best friend opioid misuse                           | 2,826    | 0.192       | 0.394     | 0          | 1          |
| Severely injured  | 2,826    | 0.138       | 0.345     | 0          | 1          |
| Any best friend severely injured                        | 2,826    | 0.161       | 0.368     | 0          | 1          |
| Smoking   | 2,817    | 0.697       | 0.460     | 0          | 1          |
| Drunk   | 2,112    | 0.705       | 0.456     | 0          | 1          |
| Unprotected sex   | 2,418    | 0.460       | 0.498     | 0          | 1          |
| Opioid addiction  | 2,521    | 0.035       | 0.184     | 0          | 1          |
| Current self-perceived health status                    | 2,826    | 2.731       | 0.918     | 0          | 4          |
| Stimulants misuse                                       | 2,826    | 0.080       | 0.272     | 0          | 1          |
| Additional outcomes (as of 2021)                        |          |             |           |            |            |
| Death due to poisoning from toxic substances or suicide | 2,826    | 0.007       | 0.081     | 0          | 1          |
| Death due to medical factors                            | 2,826    | 0.013       | 0.112     | 0          | 1          |
| Characteristics (Wave IV)                               |          |             |           |            |            |
| Female  | 2,826    | 0.523       | 0.500     | 0          | 1          |
| College   | 2,826    | 0.370       | 0.483     | 0          | 1          |
| African American  | 2,826    | 0.123       | 0.329     | 0          | 1          |
| Hispanic  | 2,826    | 0.081       | 0.273     | 0          | 1          |
| Age   | 2,826    | 28.63       | 1.771     | 25         | 34         |
| Ever diagn. depressed                                   | 2,826    | 0.150       | 0.357     | 0          | 1          |
| Ever diagn. post-traumatic stress                       | 2,826    | 0.029       | 0.169     | 0          | 1          |
| Ever diagn. anxiety                                     | 2,826    | 0.126       | 0.332     | 0          | 1          |
| Characteristics (Wave I)                                |          |             |           |            |            |
| Cigarettes avail. at parental home in WI                | 2,826    | 0.312       | 0.463     | 0          | 1          |
| Alcohol avail. at parental home in WI                   | 2,826    | 0.312       | 0.463     | 0          | 1          |

**Note:** Characteristics of individuals in the Add Health regression sample. Survey weights are used.

opioid addiction, while around 0.7 percent died due to poisoning from toxic substances or suicide by 2021.

Around 19 percent of the individuals have at least one high-school best friend who reported misusing opioids and 16 percent have at least one high-school best friend who suffered a serious injury in the previous year. Wave IV also contains information on additional socio-demographic characteristics for the respondents, such as race, occupation, and whether they completed college. There is also information on whether the respondents were ever diagnosed with depression, post-traumatic stress disorder, anxiety or panic disorder, and whether they were ever at risk under the influence of a drug. The survey in Wave IV also elicited information on the Big 5 personality traits (extraversion, agreeableness, openness, conscientiousness, and neuroticism) and a measure of risk aversion.<sup>11</sup> We also define additional measures of risky behavior based on

Wave IV responses: smoking, defined as ever having smoked an entire cigarette; alcohol use, defined as being

<sup>11</sup>The question on risk aversion was “How much do you agree with the statement about you as you generally are now, not as you wish to be in the future? ‘I like to take risks’”, and the respondents could answer on a five-point scale ranging from strongly agree to strongly disagree.

drunk or very high on alcohol at least once in the past year; and unprotected sex, defined as not having used any type of condom as a method of disease prevention in the past year.

Geographical information, such as respondents’ state, county, or Census tract of residence, is anonymized in Add Health. Hence, beyond identifying whether two individuals are from the same location, it is impossible to merge it with local-level information from external sources. However, after the conclusion of the most recent wave, conducted from 2016 to 2018, Add Health created several local-level descriptors on outcomes that are likely to interact with the effect of peers on opioid misuse. As a result, while the respondents’ county of residence is anonymous, it is possible to obtain the average characteristics of the state or county in which they live. In particular, Add Health provides state-level information on opioid dispensing rates (number of prescriptions per 100 persons) in 2016 and the number of months between the interview date and the implementation of naloxone access laws.<sup>12</sup> It also provides county-level measures on the average number of poor mental health days per month, the number of primary care physicians per 100,000 population at the county level, and information on the number of mental health facilities within 20 miles of the respondent’s residence.<sup>13</sup> Finally, a county-level social capital index, based on the methodology of [Rupasingha et al. \[2006\]](#), is also provided for 2014.<sup>14</sup>

### 3 Empirical strategy

Estimating peer effects entails several empirical challenges related to homophily, the reflection problem, and correlated effects [[Manski, 1993, 2000, Angrist, 2014](#)]. To address these, the empirical strategy combines the use of a rich data structure with an instrumental variables (IV) approach.

The first challenge, homophily, arises because friendships form endogenously—individuals tend to select friends who are similar to themselves. Consequently, observed behavioral similarities may reflect this endogenous friendship selection rather than true peer influence. Homophily is mitigated by focusing on

<sup>12</sup>State-level opioid dispensing rates provided by Add Health are from the Centers for Disease Control and Prevention (2020), <https://www.cdc.gov/drugoverdose/rxrate-maps/county2016.html>. The implementation of naloxone access laws are based on [Lee et al. \[2021\]](#). These laws provided civil or criminal immunity to licensed health care clinicians or laypersons for administering opioid antagonists, such as naloxone hydrochloride, to reverse overdoses.

<sup>13</sup>County-level health outcomes provided by AddHealth are taken from the 2020 County Health Rankings & Roadmaps, <https://www.countyhealthrankings.org/>.

<sup>14</sup>The index is based on the following establishments in each county: (a) civic organizations; (b) bowling centers; (c) golf clubs; (d) fitness centers; (e) sports organizations; (f) religious organizations; (g) political organizations; (h) labor organizations; (i) business organizations; and (j) professional organizations.

friendships formed during high school, in particular best friends, who are likely to maintain contact after high school graduation. This approach excludes new friendships formed in adulthood, which may involve individuals who misuse opioids or provide access to them after opioid misuse has begun. Additionally, by estimating an intention-to-treat effect without conditioning on current friendships, potential bias from these later endogenous relationships is avoided.<sup>15</sup>

The second challenge is the reflection problem, where simultaneity of behaviors between peers complicates disentangling the direction of influence. This challenge is addressed by employing a valid instrument—best friends’ serious injuries—which provides exogenous variation in peer opioid misuse. The rationale is that best friends who were seriously injured were prescribed opioids, which in turn led to opioid misuse.

The third challenge concerns correlated effects, which occur when individuals and their peers share unobserved environmental or contextual factors—such as local conditions or policies—that influence behavior. This issue is addressed by including fixed effects at various levels (e.g., state, school). Additionally, the influence of non-best friends residing in the same county is examined to further account for shared local factors, and these non-best friends are found to have no significant effect. This is complemented by a placebo analysis using smoking, being drunk, and having unprotected sex as alternative outcomes to rule out confounding correlated environmental influences.

Because the identification strategy relies on an instrumental variable, it is essential to ensure that both the exclusion restriction and conditional independence assumptions are satisfied. The exclusion restriction requires that a best friend’s serious injury affects the individual’s opioid misuse only through its impact on the best friend’s misuse, and not through any direct or alternative channel. This assumption could be violated if, for example, the individual and their best friend were injured in the same incident (e.g., a joint accident), or if the individual misused opioids due to emotional distress caused by the friend’s injury. To mitigate these concerns, the baseline specification controls for whether the individual experienced a serious injury and whether they have ever been diagnosed with depression, anxiety, or post-traumatic stress disorder. As a direct test of the exclusion restriction, own injuries are regressed on best friend injuries. The lack of a significant relationship confirms that simultaneous injuries—such as those from a shared accident—are

<sup>15</sup>In Sections 4.1 and 4.3, the sample is restricted to individuals who reported never using drugs in Wave I, ensuring that any potential peer influence on opioid misuse during high school is excluded.

unlikely, supporting the validity of the instrument.

Moreover, after controlling for a serious injury, individuals with and without a serious injury should have the same propensity to misuse opioids, supporting the conditional independence assumption. Conditional independence requires that, after controlling for observed characteristics and applying sample restrictions, the instrument—best friend’s serious injury—is independent of unobserved factors that directly influence the individual’s opioid misuse. In practical terms, this means that, conditional on these controls, individuals with and without a serious injury should have the same underlying propensity to misuse opioids, so that the variation in peer opioid misuse induced by the instrument is effectively random with respect to other determinants of the individual’s behavior. To support this assumption, the analysis applies several sample restrictions: i) individuals with prior illegal drug or non-prescribed painkiller use are excluded; ii) injuries related to vehicle accidents are removed; iii) individuals and their best friends identified as risk takers or engaging in risky behavior under the influence of drugs are excluded; and iv) those with recent severe injuries themselves are excluded. These restrictions help minimize confounding from unobserved traits correlated with both injury status and opioid misuse. Additionally, the placebo tests using other risky behaviors rule out that the results merely reflect a general tendency toward risky behavior among friends, providing further support for the identification strategy.

To implement the empirical strategy, the analysis first estimates a reduced-form model linking individuals’ opioid misuse directly to their best friends’ serious injuries, followed by an instrumental variables approach that uses these injuries as an instrument for peer opioid misuse.

Consider first the “reduced-form” regression that directly regresses own opioid misuse on best friends’ serious injuries:

$$\text{Opioid misuse}_{is} = \beta_1(\text{any best friend serious injury})_{is} + \beta_2(\text{serious injury})_{is} + \beta_3 X_{is} + \eta_s + u_{is}, \quad (1)$$

where  $i$  stands for the individual and  $s$  for the state of residence or school. The vector  $X_{is}$  includes socio-demographic characteristics, such as age, gender, education (with or without a college degree), and race; an indicator for having ever been diagnosed with depression, post-traumatic stress, or anxiety; and an indicator for the availability of cigarettes or alcohol in the parental home while in high school. Standard errors are

clustered using the school identifier, as recommended by the Add Health guidelines, which specify that schools are the appropriate sampling units for clustering. Longitudinal survey weights are applied to address any possible attrition, ensuring that the analysis accounts for the study’s sampling design and remains representative.

Next, consider the benchmark empirical model, which employs best friends’ serious injuries as an instrument for best friends’ opioid misuse. This instrumental variables framework includes a second-stage equation, a first-stage regression, and an exclusion restriction:

$$\begin{aligned} \text{Opioid misuse}_{is} = & \beta_1(\widehat{\text{any best friend opioid misuse}})_{is} + \beta_2(\text{serious injury})_{is} \\ & + \beta_3 X_{is} + \eta_s + u_{is}, \end{aligned} \tag{2}$$

$$\begin{aligned} \text{Any best friend opioid misuse}_{is} = & \gamma_1(\text{any best friend serious injury})_{is} + \gamma_2(\text{serious injury})_{is} \\ & + \gamma_3 X_{is} + \mu_s + e_{is}, \end{aligned} \tag{3}$$

and

$$\text{Cov}(\text{any best friend serious injury}_{is}, u_{is} \mid X_{is}, \eta_s) = 0, \tag{4}$$

The empirical specification instruments whether any best friend reports opioid misuse with whether any best friend suffered a severe injury in the previous year. The outcome variable in the benchmark specification is binary and refers to opioid misuse at any point in time prior to the Wave IV interview. Additional exercises restrict the reference period to the previous 12 months and also consider varying intensities of opioid misuse—see Section 4.3.

The benchmark OLS and 2SLS specifications include Wave III state fixed effects to control for policies that may affect the availability of opioids at the state level.<sup>16</sup> In robustness exercises, school fixed effects or a combination of school and state fixed effects are used and all results hold.

The robustness of the benchmark OLS and 2SLS estimates is checked by including an extensive list of additional individual controls; i.e., the Big 5 personality traits (extraversion, neuroticism, agreeableness, openness, conscientiousness), risk aversion, occupational dummies, the availability of drugs in the parental

<sup>16</sup>Wave III instead of Wave IV state fixed effects are used since the latter are endogenous (individuals in Wave IV may choose the state of residence based on the availability of opioids).

home while in high school, and additional family-of-origin controls (maternal education, household gross parental income during high school, and living with both parents during high school). One robustness exercise includes (exogenous) peer characteristics, namely, the fraction of best friends who are college graduates, Hispanic, African American, and the fraction who had cigarettes/alcohol/drugs available in the parental home while in high school. Other robustness exercises consider different measures of peer influence (percentage, number or dyads of best friends who misuse opioids instead of any best friend misusing opioids). The stability of peer effect estimates across these specifications further reinforces the plausibility of conditional independence.

## 4 Results

Prior to implementing the 2SLS estimation, the reduced-form OLS regression specified in equation 1 is estimated. The results are presented in Table 3.

Table 3, column 1 shows the estimated effect of best friends' serious injuries on own opioid misuse without any controls. There is a positive and statistically significant effect. The estimated effect does not change much when own severe injury (column 2) is added and is reduced in size as soon as state fixed effects (column 3) are included.

Table 3, column 1 shows the estimated causal effect of best friends' serious injuries on an individual's own opioid misuse without any controls. The effect is positive and statistically significant. When controlling for the individual's own severe injury in column 2, the estimated peer effect remains largely unchanged, indicating it is not driven by personal injury experiences. Including state fixed effects in column 3 leads to a modest reduction in the coefficient, but the effect remains statistically significant and similar in magnitude. The estimated peer effect decreases only slightly when demographic characteristics (column 4) and mental health conditions such as depression, post-traumatic stress, or anxiety (column 5) are controlled for. Including these mental health controls helps address potential confounding from shared traumatic experiences or emotional distress that might independently influence opioid misuse, ensuring the peer effect is not simply capturing these indirect pathways. Additionally, controlling for the availability of cigarettes or alcohol in the parental home during high school (column 6) rules out early household exposure to substances as a confounder. The

stability of the estimates across all specifications supports the conclusion that friends' serious injuries exert a direct and robust causal influence on opioid misuse.

To quantify the effect, the coefficient in Table 3, column 6, row 1 implies that having any best friend who has experienced a serious injury increases the probability of the respondent misusing opioids by 7.3 percentage points. This effect is substantial, considering that the average incidence of ever misusing opioids is 17 percent (Table 2). For comparison, experiencing a serious injury oneself increases the probability of opioid misuse by 9.1 percentage points (Table 3, column 6, row 3). This effect is substantial, about 80 percent of the estimated effect of experiencing a severe injury oneself, suggesting a large role for peer diffusion in opioid misuse.

Table 3: Peer effects on opioid misuse: reduced-form (OLS)

|  | Dep. var.: Prob(Opioid misuse) |                     |                     |                      |                      |                      |
|--|--------------------------------|---------------------|---------------------|----------------------|----------------------|----------------------|
|  | (1)                            | (2)                 | (3)                 | (4)                  | (5)                  | (6)                  |
| Any best friend severely injured         | 0.089***<br>(0.031)            | 0.095***<br>(0.031) | 0.079***<br>(0.028) | 0.077***<br>(0.028)  | 0.073**<br>(0.029)   | 0.073**<br>(0.029)   |
| Severely injured                         |                                | 0.123***<br>(0.031) | 0.111***<br>(0.031) | 0.099***<br>(0.031)  | 0.091***<br>(0.030)  | 0.091***<br>(0.030)  |
| College                                  |                                |                     |                     | -0.023<br>(0.021)    | -0.018<br>(0.021)    | -0.018<br>(0.023)    |
| Female                                   |                                |                     |                     | -0.035*<br>(0.019)   | -0.053***<br>(0.018) | -0.052***<br>(0.019) |
| Age                                      |                                |                     |                     | -0.017***<br>(0.005) | -0.017***<br>(0.005) | -0.018***<br>(0.006) |
| Hispanic                                 |                                |                     |                     | -0.027<br>(0.037)    | -0.027<br>(0.034)    | -0.025<br>(0.034)    |
| African American                         |                                |                     |                     | -0.060**<br>(0.026)  | -0.051**<br>(0.025)  | -0.048*<br>(0.025)   |
| Ever diagn. depressed                    |                                |                     |                     |                      | 0.098**<br>(0.040)   | 0.097**<br>(0.039)   |
| Ever diagn. post-traumatic stress        |                                |                     |                     |                      | 0.073<br>(0.069)     | 0.073<br>(0.069)     |
| Ever diagn. anxiety                      |                                |                     |                     |                      | 0.042<br>(0.043)     | 0.042<br>(0.043)     |
| Cigarettes avail. at parental home in WI |                                |                     |                     |                      |                      | 0.021<br>(0.029)     |
| Alcohol avail. at parental home in WI    |                                |                     |                     |                      |                      | 0.040<br>(0.024)     |
| Observations                             | 2,846                          | 2,846               | 2,843               | 2,830                | 2,830                | 2,826                |
| State FE                                 | No                             | No                  | Yes                 | Yes                  | Yes                  | Yes                  |
| Mean                                     | 0.168                          | 0.168               | 0.168               | 0.169                | 0.169                | 0.169                |

**Note:** The estimated coefficients of equation 1 with different sets of control variables and fixed effects (columns 1-6). Opioid misuse is defined as using prescription painkillers (e.g., Vicodin, OxyContin, Percocet, Demerol, Percodan, Tylenol with codeine) without a prescription, or taking them in larger amounts, more often, longer than prescribed, or solely for the feeling or experience they cause. Severely injured refers to experiencing any serious injury such as broken bones, cuts or lacerations, burns, torn muscles, tendons or ligaments, or other injuries that interfered with daily activities. Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \*p<0.10; \*\*p<0.05; \*\*\*p<0.01.

Building on the reduced-form evidence, the analysis turns to the instrumental variables approach to



estimate the peer effect. Table 4 reports the estimates of the second-stage regression equation (2). The first-stage results are reported in Table 5. The estimated peer effect on opioid misuse is positive and statistically significant across all specifications. Controlling for own severe injury, state fixed effects, demographic factors, mental health diagnoses (depression, post-traumatic stress, anxiety), and parental availability of cigarettes and alcohol in high school leads to only modest changes in the coefficient. This robustness indicates that the peer effect is not driven by these confounding factors or shared environmental influences. Overall, the IV estimates confirm a strong and significant peer effect on opioid misuse, aligning with the reduced-form results and reinforcing the validity of the identification strategy.

Table 4: Peer effects on opioid misuse-2SLS

|   | Dep. var.: Prob(Opioid misuse) |                     |                     |                     |                     |                     |
|---|--------------------------------|---------------------|---------------------|---------------------|---------------------|---------------------|
|   | (1)                            | (2)                 | (3)                 | (4)                 | (5)                 | (6)                 |
| Any best friend opioid misuse           | 0.576**<br>(0.227)             | 0.613***<br>(0.225) | 0.519**<br>(0.203)  | 0.495**<br>(0.196)  | 0.471**<br>(0.198)  | 0.472**<br>(0.196)  |
| Severely injured                        |                                | 0.132***<br>(0.029) | 0.124***<br>(0.029) | 0.115***<br>(0.029) | 0.106***<br>(0.029) | 0.106***<br>(0.029) |
| College                                 |                                |                     |                     | -0.038*<br>(0.023)  | -0.033<br>(0.022)   | -0.032<br>(0.023)   |
| Female                                  |                                |                     |                     | -0.024<br>(0.020)   | -0.040**<br>(0.020) | -0.040**<br>(0.020) |
| Age                                     |                                |                     |                     | -0.009<br>(0.007)   | -0.008<br>(0.007)   | -0.009<br>(0.007)   |
| Hispanic                                |                                |                     |                     | 0.000<br>(0.042)    | -0.002<br>(0.040)   | -0.000<br>(0.039)   |
| African American                        |                                |                     |                     | -0.002<br>(0.037)   | 0.002<br>(0.036)    | 0.005<br>(0.036)    |
| Ever diagn. depressed                   |                                |                     |                     |                     | 0.071*<br>(0.042)   | 0.070*<br>(0.042)   |
| Ever diagn. post-traumatic stress       |                                |                     |                     |                     | 0.100<br>(0.064)    | 0.099<br>(0.063)    |
| Ever diagn. anxiety                     |                                |                     |                     |                     | 0.041<br>(0.044)    | 0.042<br>(0.045)    |
| Cigarette avail. in parental home in WI |                                |                     |                     |                     |                     | 0.021<br>(0.031)    |
| Alcohol avail. in parental home in WI   |                                |                     |                     |                     |                     | 0.023<br>(0.028)    |
| Observations                            | 2,846                          | 2,846               | 2,843               | 2,830               | 2,830               | 2,826               |
| State FE                                | No                             | No                  | Yes                 | Yes                 | Yes                 | Yes                 |
| Kleibergen-Paap Wald rk F statistic     | 17.43                          | 17.21               | 16.90               | 18.58               | 18.27               | 18.44               |

**Note:** The estimated coefficients of equation 2 and the Kleibergen-Paap Wald rk F-statistic of equation 3 with different sets of control variables and fixed effects (columns 1-6). The estimates in column 6 correspond to the benchmark specification. Opioid misuse is defined as using prescription painkillers (e.g., Vicodin, OxyContin, Percocet, Demerol, Percodan, Tylenol with codeine) without a prescription, or taking them in larger amounts, more often, longer than prescribed, or solely for the feeling or experience they cause. Severely injured refers to experiencing any serious injury such as broken bones, cuts or lacerations, burns, torn muscles, tendons or ligaments, or other injuries that interfered with daily activities. Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \*p<0.10; \*\*p<0.05; \*\*\*p<0.01.

In all specifications, the Kleibergen-Paap Wald rk F-statistic of the first stage is above 10 (last row in Table 4), indicating that the instrument is not weak. To understand the economic significance of the

Table 5: First-stage regression

|   | Dep. var.: Prob(Any best friend opioid misuse) |
|---|--|
| Any best friend severely injured        | 0.156***<br>(0.036)                            |
| Severely injured                        | -0.033<br>(0.026)                              |
| College                                 | 0.029<br>(0.029)                               |
| Female                                  | -0.025<br>(0.024)                              |
| Age                                     | -0.018***<br>(0.007)                           |
| Hispanic                                | -0.052<br>(0.036)                              |
| African American                        | -0.113***<br>(0.028)                           |
| Ever diagn. depressed                   | 0.057*<br>(0.034)                              |
| Ever diagn. post-traumatic stress       | -0.056<br>(0.055)                              |
| Ever diagn. anxiety                     | 0.002<br>(0.031)                               |
| Cigarette avail. in parental home in WI | 0.001<br>(0.022)                               |
| Alcohol avail. in parental home in WI   | 0.036<br>(0.027)                               |
| Observations                            | 2,826  |
| State FE                                | Yes  |

**Note:** The estimated coefficients of equation 3 for the benchmark specification. Opioid misuse is defined as using prescription painkillers (e.g., Vicodin, OxyContin, Percocet, Demerol, Percodan, Tylenol with codeine) without a prescription, or taking them in larger amounts, more often, longer than prescribed, or solely for the feeling or experience they cause. Severely injured refers to experiencing any serious injury such as broken bones, cuts or lacerations, burns, torn muscles, tendons or ligaments, or other injuries that interfered with daily activities. Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \* $p < 0.10$ ; \*\* $p < 0.05$ ; \*\*\* $p < 0.01$ .

results, consider both the first-stage and second-stage coefficients. As Table 5 shows, the coefficient of the first stage is 0.156, implying that if any best friend has a serious injury, the probability that they will misuse opioids increases by 15.6 pp.<sup>17</sup> The coefficient in the second stage is 0.472 (Table 4, column 6). Therefore, if any best friend has a serious injury, the probability of the respondent misusing opioids increases by  $0.156 \times 0.472 = 0.0736$ , or 7.36 pp. Another way to interpret the magnitude is to note that the estimated effect in the second stage is conditional on having a friend who experienced an injury. As Table 2 shows, 16.1 percent of individuals have at least one best friend who was injured. Hence, if any best friend misuses opioids following a serious injury, the probability of the respondent misusing opioids increases by  $0.161 \times 0.472 = 0.0760$ , or 7.60 pp. Both of these estimates align closely with the reduced-form estimate reported in Table 3, column 6.<sup>18</sup>

<sup>17</sup>The baseline probability of any best friend misusing opioids is 19.2 percent (second row in Table 2). Recall that the period of reference is 2008, when the prescription of opioids was a common practice, especially in the case of a severe injury.

<sup>18</sup>Fletcher [2012] examines peer effects on alcohol use and finds that a 100 percent increase in classmates' alcohol use increases

## 4.1 Conditional independence and exclusion restriction

A key requirement of the instrumental variable approach is that the instrument satisfies both the exclusion restriction and conditional independence assumptions [Angrist and Pischke, 2009]. Conditional independence requires that a best friend’s serious injury be unrelated to unobserved factors affecting the individual’s opioid misuse. To strengthen this assumption, this section re-estimates the 2SLS (equations 2 and 3) excluding individuals who reported prior illegal drug use (Wave I) or non-prescribed painkiller use (Wave III), ensuring injury status is not endogenous to past substance use. Additional sample restrictions remove injuries related to vehicle accidents (potentially reflecting reckless behavior), exclude individuals and best friends who self-identify as risk takers or admit to risky behavior under the influence of drugs, and exclude individuals who themselves experienced a severe injury in Wave IV. These restrictions minimize confounding from shared risk environments, risky behavior, and injury endogeneity, improving the plausibility of conditional independence.

Table 6: Peer effects on opioid misuse-2SLS conditional independence

|                                     | Dep. var.: Prob(Opioid misuse)              |                                |                    |                             |
|-------------------------------------|---|--------------------------------|--------------------|-----------------------------|
|                                     | (1)   | (2)                            | (3)                | (4)                         |
| Any best friend opioid misuse       | 0.483**<br>(0.239)                          | 0.468**<br>(0.201)             | 0.449**<br>(0.208) | 0.420**<br>(0.195)          |
| Observations                        | 2,826                                       | 2,633                          | 2,598              | 2,476                       |
| Controls                            | Yes   | Yes                            | Yes                | Yes                         |
| FE                                  | State                                       | State                          | State              | State                       |
| Description                         | Severe injuries not due to vehicle accident | Not at risk under drugs by WIV | No risk lovers     | No own severe injury in WIV |
| Kleibergen-Paap Wald rk F statistic | 15.57                                       | 17.18                          | 16.60              | 19.67                       |

**Note:** The estimated coefficients of equation 2 and the Kleibergen-Paap Wald rk F-statistic of equation 3. Column 1 excludes injuries related to vehicle accidents. Columns 2 and 3 exclude individuals and peers with indicators of risky behavior. Column 4 excludes individuals who experienced a severe injury themselves. Robust standard errors in parentheses are clustered at the school level. Opioid misuse is defined as using prescription painkillers (e.g., Vicodin, OxyContin, Percocet, Demerol, Percodan, Tylenol with codeine) without a prescription, or taking them in larger amounts, more often, longer than prescribed, or solely for the feeling or experience they cause. Severely injured refers to experiencing any serious injury such as broken bones, cuts or lacerations, burns, torn muscles, tendons or ligaments, or other injuries that interfered with daily activities. Survey weights are used. \* $p < 0.10$ ; \*\* $p < 0.05$ ; \*\*\* $p < 0.01$ . The sample is restricted to individuals who had not used any illegal drugs in Wave I or any painkillers in Wave III. See Table 4, column 6 for the list of controls.

Table 6 presents the results. Column 1 estimates the benchmark specification considering only own and best friends’ severe injuries that did not involve a vehicle accident. The estimates remain robust, suggesting that the results are not driven by vehicular accidents potentially caused by risky behavior associated with the likelihood of drinking by 50 percentage points in a population where 48 percent drink alcohol. The peer effects we find are stronger (a 47 percentage points increase in a population where 17 percent misuse opioids). This can be due to differences in IV strategies. During the peak of the opioid epidemic, individuals who experienced a severe injury were almost certain to be treated with opioids, which amplified the peer effect, as the probability of misusing opioids if you get prescribed opioids after an injury was 15.6% (Table 5). In contrast, Fletcher [2012] uses the availability of alcohol at classmates’ parental homes as an instrument.

opioid misuse. Column 2 retains only individuals and best friends who either do not use drugs in Wave IV or they use drugs but report that they have never put themselves or others at risk under the influence of a drug.<sup>19</sup> In this way, serious injuries that may have occurred due to drug use are excluded. In a similar vein, column 3 excludes individuals and best friends who are risk lovers (answered strongly agree to the statement ‘I like to take risks’). The results of both exercises are extremely similar to the benchmark estimates. Last, column 4 excludes individuals who suffered a severe injury themselves in Wave IV. Also in this case, the estimated peer effects are highly statistically significant and of similar size as in the benchmark specification. Across all specifications, the estimated peer effects remain stable and statistically significant, reinforcing the credibility of the identification strategy and the plausibility of the conditional independence assumption.

Table 7: Testing the exclusion restriction: Effect of best friends’ injury on own injury

|                                  | Dep. var.: Prob(Severely injured) |                   |
|----------------------------------|-----------------------------------|-------------------|
|                                  | (1)                               | (2)               |
| Any best friend severely injured | -0.023<br>(0.027)                 | -0.034<br>(0.024) |
| Observations                     | 2,131                             | 2,118             |
| Controls                         | No                                | Yes               |
| State FE                         | Yes                               | Yes               |
| Mean                             | 0.117                             | 0.116             |

**Note:** Robust standard errors in parentheses are clustered at the school level. Severely injured refers to experiencing any serious injury such as broken bones, cuts or lacerations, burns, torn muscles, tendons or ligaments, or other injuries that interfered with daily activities. Survey weights are used. \* $p < 0.10$ ; \*\* $p < 0.05$ ; \*\*\* $p < 0.01$ . The sample is restricted to individuals who had not used any illegal drugs in Wave I or any painkillers in Wave III. See Table 4, column 6 for the list of controls.

The exclusion restriction requires that the instrument affect the individual’s opioid misuse only through the best friend’s misuse, with no direct or alternative channels. To address potential violations of the exclusion restriction, such as simultaneous injuries from the same accident or emotional distress caused by a friend’s injury, the baseline specification controls for the individual’s own serious injuries as well as diagnoses of depression, anxiety, and post-traumatic stress disorder. Moreover, Table 7 reports a direct test of the exclusion restriction, which examines whether an individual’s injury status is correlated with that of their best friend. Specifically, own injuries are regressed on best friend injuries. The absence of a statistically significant relationship suggests that simultaneous injury—such as those resulting from a shared accident—are

<sup>19</sup>More specifically, those who answered “never” to the following question in Wave IV are kept: “How often have you been under the influence of your favorite drug when you could have gotten yourself or others hurt, or put yourself or others at risk, including unprotected sex?” Within the context of the Add Health questions, drugs here exclude misuse of prescription opioids.

unlikely. This supports the claim that a best friend’s serious injury affects the individual’s opioid misuse only through its impact on the friend’s behavior, not through any direct or alternative channel. These results reinforce the validity of the exclusion restriction and lend credibility to the causal interpretation of the estimated peer effects.

## 4.2 Placebo exercises

The reduced-form equation can also be used to exclude the possibility that the benchmark specification captures some general risky behavior among friends rather than a pure peer effect. To this end, equation 1 is estimated with the probability of smoking, being drunk, and having unprotected sex rather than the probability of opioid misuse as outcome variables. As Table 8, columns 1-3, show, the estimated coefficients in the placebo regressions are essentially null, further supporting the interpretation of the estimated effect in the benchmark specification as a genuine peer effect. These placebo tests help rule out the possibility that the results are driven by shared environmental factors or a general propensity for risky behavior among peers, rather than the specific influence of exposure to opioid-related injuries.

Table 8: Placebo exercises-OLS

| Dep. var.:                       | Prob(Smoking)<br>(1) | Prob(Drunk)<br>(2) | Prob(Unprotected sex)<br>(3) |
|----------------------------------|----------------------|--------------------|------------------------------|
| Any best friend severely injured | 0.004<br>(0.031)     | 0.023<br>(0.037)   | 0.036<br>(0.036)             |
| Severely injured                 | 0.029<br>(0.039)     | 0.009<br>(0.042)   | -0.019<br>(0.041)            |
| Observations                     | 2,817                | 2,111              | 2,418                        |
| State FE                         | Yes                  | Yes                | Yes                          |
| Controls                         | Yes                  | Yes                | Yes                          |
| Mean                             | 0.697                | 0.705              | 0.460                        |

**Note:** The estimated coefficients of equation 1 for placebo outcomes. The dependent variables are: in column (1), the probability of smoking, defined as ever having smoked an entire cigarette; in column (2), the probability of being drunk or very high on alcohol at least once in the past year; and in column (3), the probability of having had unprotected sex in the past year, defined as not having used any type of condom as a method of disease prevention. Severely injured refers to experiencing any serious injury such as broken bones, cuts or lacerations, burns, torn muscles, tendons or ligaments, or other injuries that interfered with daily activities. Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \*p<0.10; \*\*p<0.05; \*\*\*p<0.01. See Table 3, column 6 for the list of controls.

### 4.3 Timing and intensity of opioid misuse

The benchmark model uses as the outcome variable whether any opioid misuse ever occurred as reported in Wave IV. This may give rise to two potential issues. First, there is a potential time discrepancy between the outcome and the instrumental variable as severe injuries reported in Wave IV refer to the past 12 months. Second, the outcome variable does not distinguish between occasional and more intensive opioid misuse. This section addresses both issues by using a different question from Add Health that specifies the intensity of painkillers misuse in the past 12 months.<sup>20</sup> Individuals with previous experience with illegal drugs or non-prescribed painkillers are again excluded from the analysis as in Section 4.1.

Table 9: Peer effects on opioid misuse-2SLS by timing and intensity of misuse

| Intensity                           | Dep. var.: Prob(Opioid misuse) |                                  |  |  |   |
|-------------------------------------|--------------------------------|----------------------------------|--|--|---|
|                                     | Ever misused                   | Any misuse in the last 12 months | Once per month or more in the last 12 months | 2-3 days per month or more in the last 12 months | 1-2 days per week or more in the last 12 months |
|                                     | (1)                            | (2)                              | (3)  | (4)  | (5)   |
| Any best friend opioid misuse       | 0.505***<br>(0.174)            | 0.314**<br>(0.124)               | 0.291**<br>(0.118)                           | 0.283**<br>(0.112)                               | 0.258**<br>(0.123)                              |
| Observations                        | 2,118                          | 2,010                            | 1,985  | 1,967  | 1,950   |
| Controls                            | Yes                            | Yes                              | Yes  | Yes  | Yes   |
| State FE                            | Yes                            | Yes                              | Yes  | Yes  | Yes   |
| Mean                                | 0.110                          | 0.0584                           | 0.0419                                       | 0.0311   | 0.0152  |
| Kleibergen-Paap Wald rk F statistic | 13.46                          | 13.77                            | 13.54  | 12.80  | 10.65   |

**Note:** The estimated coefficients of equation 2 and the Kleibergen-Paap Wald rk F-statistic of equation 3 for different intensities of opioid misuse in the previous 12 months. Opioid misuse is defined as using prescription painkillers (e.g., Vicodin, OxyContin, Percocet, Demerol, Percodan, Tylenol with codeine) without a prescription, or taking them in larger amounts, more often, longer than prescribed, or solely for the feeling or experience they cause. Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \*p<0.10; \*\*p<0.05; \*\*\*p<0.01. The sample is restricted to individuals who had not used any illegal drugs in Wave I or any painkillers in Wave III. See Table 4, column 6 for the list of controls.

Table 9 presents the results using this sample restriction for opioid misuse across different time frames (ever or past 12 months) and with progressively increasing levels of misuse intensity.<sup>21</sup> The coefficient for the peer effect remains statistically significant and comparable in magnitude to the benchmark specification when the sample is limited to individuals who had not used illegal drugs in Wave I or painkillers in Wave III (column 1). This holds true even when considering opioid misuse only within the past 12 months.<sup>22</sup>

<sup>20</sup>The question is: “During the past 12 months, on how many days did you use painkillers?: none, 1 or 2 days, once a month or less, 2 or 3 days a month, 1 or 2 days a week, 3 to 5 days a week, every day or almost every day.”

<sup>21</sup>It is important to note that only the misuse intensity of the affected individual is adjusted, while the peers’ misuse intensity remains unchanged. Peers are not excluded based on their level of misuse, ensuring that the peer group composition remains consistent with the benchmark analysis. Consequently, the first-stage relationship, which estimates the impact of a peer’s injury on opioid misuse among peers, remains unaffected by the exclusion of individuals with lower-intensity misuse.

<sup>22</sup>All specifications, robustness checks, and group heterogeneity analyses hold when the analysis is repeated using this variable and sample restriction as a benchmark. However, some estimates become less precise due to the smaller sample size.

Peer effects remain significant as the intensity of opioid misuse increases (columns 3 and 4) and persist even at a relatively high level of misuse (1-2 days per week, as shown in column 5). The coefficients decrease in magnitude as misuse intensity rises, reflecting the lower prevalence of more intensive misuse within the population, as indicated by the mean. Overall, these findings indicate that peer effects on opioid misuse are robust across different definitions of misuse timing and intensity, and extend beyond occasional misuse to more serious levels.

#### 4.4 Robustness checks

A battery of additional exercises are run to check the robustness of our benchmark estimates. Figure 1 and Table A1 report the results. The first exercise (E1 in Figure 1) introduces school instead of state fixed effects and the estimates continue to be statistically significant but of slightly smaller size. The second exercise (E2 in Figure 1) includes both state and school fixed effects. Although this specification is demanding (it includes 174 fixed effects), the coefficient continues to be statistically significant and of slightly smaller size than the benchmark estimate.




Figure 1: Peer effects on opioid misuse-2SLS robustness

**Note:** The estimated coefficients and 90% confidence intervals of equation 2 with different sets of fixed effects, control variables, and sample restrictions. See Table A1 for the full set of estimates.

Exercises E3 and E4 control for whether drugs were available in the parental home in Wave I and for

exogenous/pre-determined friend characteristics. The results are very similar to the benchmark estimate. The findings are also robust to the inclusion of the Big 5 personality traits and a risk aversion measure (exercise E5) and to the inclusion of occupational dummies (exercise E6).<sup>23</sup> Last, exercise E7 controls for maternal education, household income, and household structure in Wave I. These variables are available for a smaller sample, but the peer effects remain significant.

Additionally, Table A2 reports estimates with different ways of measuring peer influence. In particular, while the benchmark regressor is whether any best friend misused opioids, column 2 shows the results with the percentage of best friends misusing opioids (0, 50% or 100%), column 3 with the number of best friends misusing opioids (0, 1 or 2) and columns 4 using dyads of best friends (i.e., considering the opioid misuse of each best friend of the respondent separately). All estimates are perfectly in line with the benchmark results.

## 5 Heterogeneous effects and mechanisms

There is substantial heterogeneity in opioid misuse by socioeconomic characteristics. In particular, opioid misuse is more common among less educated individuals and among non-Hispanic whites. As Table A3 shows, in Wave IV of Add Health, less than 15 percent of college graduates report ever using opioids, whereas the number for non-college graduates is 18.2 percent.<sup>24</sup> Moreover, 19.3 percent of non-Hispanic whites report ever using opioids in Wave IV, whereas the number is just 6.3 percent for other races.

Some individuals are more prone to the influence of peers than others. This is examined by focusing on education and race and distinguishing between college and non-college graduates and between non-Hispanic whites and others. Table 10 presents the results. Peer effects in the second stage are significant and large for those without a college degree (column 2) but not among those with a college degree (column 1). There is no significant difference in the peer effect by race (columns 3 and 4).<sup>25</sup> The F-statistic of the first stage is close to or above 10 for all groups (last row of Table 10).

<sup>23</sup>Previous research has shown that personality traits are an important determinant of other health-related outcomes such as bulimia. See Ham et al. [2021]. The current analysis shows that risk aversion and conscientiousness decrease the probability of opioid misuse while openness increases it.

<sup>24</sup>The educational gap in opioid misuse is higher in the NSDUH. Between 2005 and 2019, among individuals between ages 26 and 34, 4.7 percent of individuals with a college degree and 7.8 percent of those without a college degree reported misusing opioids during the last 12 months.

<sup>25</sup>Due to the small sample size, only two race categories, Non-Hispanic Whites and Others, are considered.



Table 10: Peer effects on opioid misuse-2SLS by education and race

|                                     | Dep. var.: Prob(Opioid misuse) |                    |                    |                  |
|-------------------------------------|--------------------------------|--------------------|--------------------|------------------|
|                                     | By education                   |                    | By race            |                  |
|                                     | College                        | Non-College        | Non-Hispanic white | Others           |
|                                     | (1)                            | (2)                | (3)                | (4)              |
| Any best friend opioid misuse       | 0.206<br>(0.300)               | 0.597**<br>(0.238) | 0.395*<br>(0.223)  | 0.691<br>(0.485) |
| Observations                        | 1,072                          | 1,747              | 1,726              | 1,095            |
| Controls                            | Yes                            | Yes                | Yes                | Yes              |
| State FE                            | Yes                            | Yes                | Yes                | Yes              |
| Mean                                | 0.127                          | 0.165              | 0.189              | 0.0728           |
| Kleibergen-Paap Wald rk F statistic | 9.829                          | 12.42              | 12.31              | 9.177            |

**Note:** The estimated coefficients of equation 2 and the Kleibergen-Paap Wald rk F-statistic of equation 3 for different educational groups (columns 1 and 2) and for different racial groups (columns 3 and 4). Opioid misuse is defined as using prescription painkillers (e.g., Vicodin, OxyContin, Percocet, Demerol, Percodan, Tylenol with codeine) without a prescription, or taking them in larger amounts, more often, longer than prescribed, or solely for the feeling or experience they cause. Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \*p<0.10; \*\*p<0.05; \*\*\*p<0.01. See Table 4, column 6 for the list of controls.

Peer effects could arise due to information sharing regarding the efficacy of opioids or through the direct provision of opioids. To shed light on the underlying mechanism, the type, duration and similarity of friendship is considered as well as the geographical proximity between the respondents and their best friends. Figure 2 and Table A4 report the results.

First, peer effects are statistically significant only among best friends (first female and male nomination). Other friends (second to fifth female and male nominations) do not exert any statistically significant effect on opioid misuse. Second, the duration of the friendship and the similarity of friends matters. The peer effect stems exclusively from best friends nominated both in Wave I and Wave II. Moreover, the peer effect becomes larger when same gender best friends are considered.

Geographical proximity is also crucial. For around half of the cases, the county of residence of the respondents and of their best friends in Wave III coincides. Therefore equation 2 is re-estimated considering best friends residing in the same county as the respondents and in a different county. The estimates show that peer effects arise exclusively from best friends in the same county, highlighting the key role of geographical proximity.<sup>26,27</sup> Note that the peer effect from non-best friends residing in the same county as the respondent

<sup>26</sup>The results are very similar with the inclusion of county or school fixed effects instead of state fixed effects.

<sup>27</sup>The peer effects are stronger for non-college graduates, who are less likely to move. The fraction of college graduates is 37 percent in the overall sample and only 32 percent in the restricted sample with same county best friends. However, if we re-run the same county best friends regression for college and non-college separately, the peer effect is more substantial for both education groups.




Figure 2: Peer effects on opioid misuse-by type of friends

**Note:** The estimated coefficients and 90% confidence intervals of equation 2 by type of friends. Estimates for best (first nominated) friends, other (second, third, fourth or fifth nominated) friends; for best friends nominated both in Wave I and Wave II, for best friends nominated only in Wave I, and for best friends of the same gender as the respondent; for best friends residing in the same or different county as the respondent, for other friends residing in the same or different county as the respondent. See Table A4 for the full set of estimates.

is null. This is reassuring as it rules out the possibility that the estimated peer effects are subject to correlated effects (e.g., a common shock that affects everyone in the county). The results in Figure 2 and Table A4 can be interpreted as supportive evidence both of the direct provision channel and information sharing among friends, who are more likely to stay in touch.

The granularity and richness of information in Add Health allows for the exploration of heterogeneity based on the characteristics of the entire network of friendship nominations. In particular, own and friends' Bonacich centrality (the centrality of an individual weighted by the centrality of those they nominate as friends), network reach (the maximum number of individuals a node can connect within the entire network), and network density (the number of actual ties in the total friendship network divided by the maximum possible density given an out-degree of 10) are considered.<sup>28</sup> On the effectiveness of interventions that target individuals who are central in their networks, see, among others, Banerjee et al. [2013]. Figure 3 and Table

<sup>28</sup>Figures A3 and A4 illustrate examples of networks with low and high centrality; and low and high density.




Figure 3: Peer effects on opioid misuse-by network characteristics

**Note:** The estimated coefficients and 90% confidence intervals of equation 2 by network characteristics. Estimates for individuals that are less or more central in the entire friendship network (Bonacich centrality below or above the mean); for individuals with a low or high reach (maximum number of individuals a node can connect within the entire friendship network is below or above the mean); and for individuals in relatively less or more dense friendship networks. See Table A5 for the full set of estimates.

A5 show the results. The peer effect is larger in dense networks and in networks with a high reach. Moreover, the analysis reveals that peer effects emanate from and influence individuals that are central or “key players” in their network.

Last, the impact of local-level factors and policies as mitigating factors on peer effects is investigated. Specifically, local variations in opioid availability, measured by opioid dispensing rates, and access to naloxone are considered. The health of the local population, assessed by the prevalence of mental health issues, and the quality of healthcare, measured by the availability of mental health facilities and primary care physicians, are also analyzed. Additionally, local differences in social capital are investigated as another potential mitigating factor.

The results (Figure 4, Table A6) show that peer effects on opioid misuse are stronger in states with high opioid dispensing rates and in states where naloxone access laws were implemented later. Delayed naloxone access may have limited harm-reduction measures, allowing stronger peer effects to develop, while high dispensing rates likely increased exposure to opioids and normalized misuse. Peer effects are also greater in counties with more poor mental health days and fewer primary-care physicians per capita, highlighting




Figure 4: Peer effects on opioid misuse: by opioid and naloxone availability; prevalence of mental health problems; social capital index; mental health facilities and physicians availability

**Note:** The estimated coefficients and 90% confidence intervals of equation 2 by local-level factors. Low/high refers to values below/above the median in the sample. Estimates for individuals who reside in states with opioid dispense rates below or above the median; for individuals who reside in states that implemented naloxone laws earlier or later; for individuals who reside in counties with an average number of poor mental health days per month below or above the median; for individuals who reside in counties with a social capital index in 2014 below or above the median; for individuals who reside in locations with less or more than 13 mental health facilities within 20 miles; for individuals who reside in counties with less or more than 70 primary-care physicians per 100,000 population. The location of residence is as of the date of the Wave V interview (that took place in the period 2016-2018). See Table A6 for the full set of estimates.

the role of unmet healthcare needs. These effects weaken when the availability of mental health facilities or primary-care physicians exceeds the sample median, underscoring the mitigating role of healthcare infrastructure. Finally, counties with low social capital exhibit stronger peer effects. This result aligns with prior evidence that links low social capital to higher rates of opioid overdose mortality (Zoorob and Salemi [2017]).

## 6 Welfare and broader implications of peer opioid misuse

A natural question is whether opioid misuse deteriorates welfare and has other broader consequences. To this end, this section analyzes the short-run and long-run implications of peer-initiated opioid misuse by considering different outcome variables. Table 11 reports the results. Column 1 shows that peer opioid

misuse is detrimental for own self-perceived health. Self-perceived health status decreases by 0.3 points ( $-0.847 \times 0.156$  from the first-stage regression) on a 5-point scale if any best friend has a serious injury. Moreover, the probability of stimulants misuse increases by 3.9 pp ( $0.251 \times 0.156$ ) in a population in which 5 percent misuse stimulants (column 2). This is not surprising as stimulants are often combined with opioids [Ahmed et al., 2022].

Table 11: Welfare and broader implications-2SLS

| Dep. Var.:                          | Current self-perceived health status | Stimulants misuse | Opioid addiction  | Death due to medical factors | Death due to poisoning from toxic substances or suicide |
|-------------------------------------|--------------------------------------|-------------------|-------------------|------------------------------|---|
|                                     | (1)                                  | (2)               | (3)               | (4)                          | (5)   |
| Any best friend opioid misuse       | -0.847*<br>(0.453)                   | 0.251*<br>(0.137) | 0.177*<br>(0.097) | -0.044<br>(0.044)            | 0.087<br>(0.080)  |
| Observations                        | 2,118                                | 2,118             | 1,958             | 2,118                        | 2,118   |
| State FE                            | Yes                                  | Yes               | Yes               | Yes                          | Yes   |
| Mean                                | 2.775                                | 0.052             | 0.015             | 0.010                        | 0.003   |
| Kleibergen-Paap Wald rk F statistic | 13.46                                | 13.46             | 10.29             | 13.46                        | 13.46   |

**Note:** The dependent variable is current self-perceived health status (5-point scale ranging from poor to excellent) in column 1; the probability of stimulants misuse in column 2; the probability of opioid addiction (continued opioid use despite mental and/or health problems due to it) in column 3; death due to medical factors (e.g., HIV, cancer, diabetes, CVD, parasitic, respiratory or digestive diseases) in column 4; and death due to poisoning from toxic substances or suicide in column 5. Reference period of opioid addiction/stimulants misuse: ever in life. Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \* $p < 0.10$ ; \*\* $p < 0.05$ ; \*\*\* $p < 0.01$ . The sample is restricted to individuals who had not used any illegal drugs in Wave I or any painkillers in Wave III. See Table 4, column 6 for the list of controls.

To examine whether peer opioid misuse has severe consequences in the long run, information on opioid addiction and causes of death is utilized. Opioid addiction refers to the continued use of opioids despite the presence of mental and/or physical health problems caused by their use. The focus here is on deaths due to causes likely related to opioid misuse, namely, poisoning from toxic substances or suicide. These cover two of the causes that Case and Deaton [2020] label as the deaths of despair: drug overdose (including alcohol overdose), suicide, and alcoholic liver disease. Deaths due to alcoholic liver disease are left out since, in the Add Health classification, they are aggregated with digestive diseases. As column 3 in Table 11 shows, peer opioid misuse leads to an increase in the probability of opioid addiction. The effect is significant, as the probability of opioid addiction increases by 2.7 pp ( $0.177 \times 0.156$ ) if any best friend has a severe injury.<sup>29</sup> Columns 4 and 5 show the effects on the deaths due to medical factors and on deaths due to causes likely to be related to opioid misuse. There is no statistically significant effect of peer opioid misuse on the probability of death due to medical factors (column 4), and the estimated coefficient is negative. Furthermore, there is a positive effect of peer opioid misuse on the likelihood of death due to poisoning from toxic substances or

<sup>29</sup>This large effect aligns with the findings of Cutler and Donahoe [2024], who use county-level data to show that spillovers between counties that are geographically close or linked through Facebook friendship networks can account for 84 to 92 percent of opioid deaths from 1990 to 2018, and are a major factor in the sustained increase in opioid-related deaths over time.

suicide (column 5). Still, as deaths are rare in this age group, the estimates are imprecise. In summary, these results suggest that peer opioid misuse can deteriorate health, increase the likelihood of opioid addiction, and potentially contribute to eventual deaths from overdose or suicide, although the estimates for such rare outcomes are imprecise.

## 7 Conclusions

Using individual-level data from Add Health and a novel identification strategy, peer effects on opioid misuse between friends are causally identified. Concerns related to endogenous friendship formation and termination are mitigated by focusing on friendships formed during high school and subsequent drug use as an adult. The spotlight is on best friends, who are likely to maintain contact after high school graduation and the estimation of an intention-to-treat effect without conditioning on current friendships. By studying opioid use at least 14 years after friendship formation and using a credible instrument (best friends’ serious injuries) the challenge of simultaneity (the reflection problem) in the estimation of peer effects is addressed. The analysis finds significant positive peer effects on opioid misuse, especially among individuals without a college degree and individuals with strong ties who are central in their network. Moreover, peer opioid misuse is likely to lead to deteriorating health and opioid addiction.

By identifying the role of peer spillovers in opioid misuse, the current study provides a more comprehensive understanding of the mechanisms behind the epidemic, complementing the supplier-driven explanations and highlighting the importance of social networks in the spread of opioid misuse. The findings have implications for the design of policies that are meant to reduce opioid dependence [Currie and Schwandt, 2021, Gupta et al., 2023]. The reformulation of OxyContin and the implementation of must-access prescription drug monitoring programs had unintended consequences, with opioid-dependent users resorting to illegal drugs including heroin [Alpert et al., 2018] and a subsequent increase in child abuse [Evans et al., 2022]. The large social multiplier that is identified suggests that policies targeted on selected individuals (e.g., those with a large social network) may be particularly effective. For example, educating juveniles about the perils of drug use via advertising campaigns on television and social media might be effective.

## References

- Effrosyni Adamopoulou and Ezgi Kaya. Young Adults Living with their Parents and the Influence of Peers. *Oxford Bulletin of Economics and Statistics*, 80(3):689–713, 2018. doi:[10.1111/obes.12198](https://doi.org/10.1111/obes.12198).
- Francesco Agostinelli, Matthias Doepke, Giuseppe Sorrenti, and Fabrizio Zilibotti. When the Great Equalizer Shuts Down: Schools, Peers, and Parents in Pandemic Times. *Journal of Public Economics*, 206:104574, 2022. doi:[10.1016/j.jpubeco.2021.104574](https://doi.org/10.1016/j.jpubeco.2021.104574).
- Saeed Ahmed, Zouina Sarfraz, and Azza Sarfraz. A Changing Epidemic and the Rise of Opioid-stimulant Co-use. *Frontiers in psychiatry*, 13:918197, 2022. doi:[10.3389/fpsyt.2022.918197](https://doi.org/10.3389/fpsyt.2022.918197).
- Abby Alpert, David Powell, and Rosalie Liccardo Pacula. Supply-Side Drug Policy in the Presence of Substitutes: Evidence from the Introduction of Abuse-Deterrent Opioids. *American Economic Journal: Economic Policy*, 10(4):1–35, 2018. doi:[10.1257/pol.20170082](https://doi.org/10.1257/pol.20170082).
- Abby Alpert, William N Evans, Ethan M J Lieber, and David Powell. Origins of the Opioid Crisis and its Enduring Impacts. *The Quarterly Journal of Economics*, 137(2):1139–1179, 2022. doi:[10.1093/qje/qjab043](https://doi.org/10.1093/qje/qjab043).
- Joshua D Angrist. The Perils of Peer Effects. *Labour Economics*, 30:98–108, 2014. doi:[10.1016/j.labeco.2014.05.008](https://doi.org/10.1016/j.labeco.2014.05.008).
- Joshua D Angrist and Jörn-Steffen Pischke. *Mostly Harmless Econometrics: An Empiricist’s Companion*. Princeton University Press, 2009.
- Abhijit Banerjee, Arun G Chandrasekhar, Esther Duflo, and Matthew O Jackson. The Diffusion of Microfinance. *Science*, 341(6144):1236498, 2013. doi:[10.1126/science.123649](https://doi.org/10.1126/science.123649).
- Michael L Barnett, Andrew R Olenski, and Anupam B Jena. Opioid-Prescribing Patterns of Emergency Physicians and Risk of Long-Term Use. *New England Journal of Medicine*, 376(7):663–673, 2017. doi:[10.1056/NEJMsa1610524](https://doi.org/10.1056/NEJMsa1610524).
- Robert Bifulco, Jason M Fletcher, and Stephen L Ross. The Effect of Classmate Characteristics on Post-secondary Outcomes: Evidence from the Add Health. *American Economic Journal: Economic Policy*, 3(1):25–53, February 2011. doi:[10.1257/pol.3.1.25](https://doi.org/10.1257/pol.3.1.25).

- Carlos Blanco, Tisha R A Wiley, Jacqueline J Lloyd, Marsha F Lopez, and Nora D Volkow. America's Opioid Crisis: The Need for an Integrated Public Health Approach. *Translational Psychiatry*, 10(167), 2020. doi:[10.1038/s41398-020-0847-1](https://doi.org/10.1038/s41398-020-0847-1).
- Lorenzo Cappellari and Konstantinos Tatsiramos. With a Little Help from my Friends? Quality of Social Networks, Job Finding and Job Match Quality. *European Economic Review*, 78:55–75, 2015. doi:[10.1016/j.euroecorev.2015.04.002](https://doi.org/10.1016/j.euroecorev.2015.04.002).
- David Card and Laura Giuliano. Peer Effects and Multiple Equilibria in the Risky Behavior of Friends. *Review of Economics and Statistics*, 95(4):1130–1149, 2013. doi:[doi.org/10.1162/REST\\_a\\_00340](https://doi.org/10.1162/REST_a_00340).
- Jacqueline A Carrico, Katharine Mahoney, Kristen M Raymond, Shannon K McWilliams, Lena M Mayes, Susan K Mikulich-Gilbertson, and Karsten Bartels. Predicting Opioid Use Following Discharge After Cesarean Delivery. *The Annals of Family Medicine*, 18(2):118–126, 2020. doi:[10.1370/afm.2493](https://doi.org/10.1370/afm.2493).
- Anne Case and Angus Deaton. *Deaths of Despair and the Future of Capitalism*. Princeton University Press: Princeton, NJ, 2020.
- Andrew E Clark and Youenn Lohéac. “It Wasn’t Me, It Was Them!” Social Influence in Risky Behavior by Adolescents. *Journal of Health Economics*, 26(4):763–784, 2007. doi:[10.1016/j.jhealeco.2006.11.005](https://doi.org/10.1016/j.jhealeco.2006.11.005).
- Wilson M Compton, Christopher M Jones, Grant T Baldwin, Frances M Harding, Carlos Blanco, and Eric M Wargo. Targeting Youth to Prevent Later Substance Use Disorder: An Underutilized Response to the US Opioid Crisis. *American Journal of Public Health*, 109(S3):S185–S189, 2019. doi:[10.2105/AJPH.2019.305020](https://doi.org/10.2105/AJPH.2019.305020).
- Janet Currie and Hannes Schwandt. The Opioid Epidemic Was Not Caused by Economic Distress But by Factors that Could Be More Rapidly Addressed. *The Annals of the American Academy of Political and Social Science*, 695(1):276–291, 2021. doi:[10.1177/00027162211033833](https://doi.org/10.1177/00027162211033833).
- Janet Currie and Esmée Zwiers. Medication of Postpartum Depression and Maternal Outcomes: Evidence from Geographic Variation in Dutch Prescribing. *Journal of Human Resources*, 2023. doi:[10.3368/jhr.1021-11986R1](https://doi.org/10.3368/jhr.1021-11986R1).



- David M Cutler and J Travis Donahoe. Thick Market Externalities and the Persistence of the Opioid Epidemic. Technical report, National Bureau of Economic Research, 2024.
- David M Cutler and Edward Glaeser. What Explains Differences in Smoking, Drinking, and Other Health-Related Behaviors? *American Economic Review*, 95(2):238–242, 2005. doi:[10.1257/000282805774670464](https://doi.org/10.1257/000282805774670464).
- David M Cutler and Edward L Glaeser. When Innovation Goes Wrong: Technological Regress and the Opioid Epidemic. *Journal of Economic Perspectives*, 35(4):171–196, 2021. doi:[10.1257/jep.35.4.171](https://doi.org/10.1257/jep.35.4.171).
- Gordon B Dahl, Katrine V Løken, and Magne Mogstad. Peer Effects in Program Participation. *American Economic Review*, 104(7):2049–2074, 2014. doi:[10.1257/aer.104.7.2049](https://doi.org/10.1257/aer.104.7.2049).
- Giacomo De Giorgi, Michele Pellizzari, and Silvia Redaelli. Identification of Social Interactions through Partially Overlapping Peer Groups. *American Economic Journal: Applied Economics*, 2(2):241–275, 2010. doi:[10.1257/app.2.2.241](https://doi.org/10.1257/app.2.2.241).
- Mathijs de Vaan and Toby Stuart. Does Intra-household Contagion Cause an Increase in Prescription Opioid Use? *American Sociological Review*, 84(4):577–608, 2019. doi:[10.1177/0003122419857797](https://doi.org/10.1177/0003122419857797).
- William N Dowd. The Effect of Untargeted Naloxone Distribution on Opioid Overdose Outcomes. *Health economics*, 32(12):2801–2818, 2023. doi:[10.1002/hec.4755](https://doi.org/10.1002/hec.4755).
- Sarah Eichmeyer and Jonathan Zhang. Pathways into Opioid Dependence: Evidence from Practice Variation in Emergency Departments. *American Economic Journal: Applied Economics*, 14(4):271–300, 2022. doi:[10.1257/app.20210048](https://doi.org/10.1257/app.20210048).
- Sarah Eichmeyer and Jonathan Zhang. Primary Care Providers’ Influence on Opioid Use and Its Adverse Consequences. *Journal of Public Economics*, 217:104784, 2023. doi:[10.1016/j.jpubeco.2022.104784](https://doi.org/10.1016/j.jpubeco.2022.104784).
- Daniel Eisenberg, Ezra Golberstein, and Janis L Whitlock. Peer Effects on Risky Behaviors: New Evidence from College Roommate Assignments. *Journal of health economics*, 33:126–138, 2014. doi:[10.1016/j.jhealeco.2013.11.006](https://doi.org/10.1016/j.jhealeco.2013.11.006).

- Mary F Evans, Matthew C Harris, and Lawrence M Kessler. The Hazards of Unwinding the Prescription Opioid Epidemic: Implications for Child Maltreatment. *American Economic Journal: Economic Policy*, 14(4):192–231, 2022. doi:[10.1257/pol.20200301](https://doi.org/10.1257/pol.20200301).
- Jesús Fernández-Villaverde, Jeremy Greenwood, and Nezih Guner. From Shame to Game in One Hundred Years: An Economic Model of the Rise in Premartial Sex and its De-Stigmatization. *Journal of the European Economic Association*, 12(1):25–61, 2014. doi:[10.1111/jeea.12043](https://doi.org/10.1111/jeea.12043).
- Amy Finkelstein, Matthew Gentzkow, Dean Li, and Heidi L Williams. What Drives Risky Prescription Opioid Use? Evidence from Migration. Technical report, National Bureau of Economic Research, 2022.
- Jason M Fletcher. Peer Influences on Adolescent Alcohol Consumption: Evidence Using an Instrumental Variables/Fixed Effect Approach. *Journal of Population Economics*, 25:1265–1286, 2012. doi:[10.1007/s00148-011-0365-9](https://doi.org/10.1007/s00148-011-0365-9).
- Sumedha Gupta, Thuy Nguyen, Patricia R Freeman, and Kosali Simon. Competitive Effects of Federal and State Opioid Restrictions: Evidence from the Controlled Substance Laws. *Journal of Health Economics*, 91:102772, 2023. doi:[10.1016/j.jhealeco.2023.102772](https://doi.org/10.1016/j.jhealeco.2023.102772).
- Gery P Guy. Vital Signs: Changes in Opioid Prescribing in the United States, 2006–2015. *MMWR. Morbidity and mortality weekly report*, 66, 2017. doi:[10.15585/mmwr.mm6626a4](https://doi.org/10.15585/mmwr.mm6626a4).
- John C Ham, Daniela Iorio, and Michelle Sovinsky. Health Outcomes, Personality Traits and Eating Disorders. *Economic Policy*, 36(105):51–76, 2021. doi:[10.1093/epolic/eiaa029](https://doi.org/10.1093/epolic/eiaa029).
- Kathleen Mullan Harris. The National Longitudinal Study of Adolescent to Adult Health (Add Health), Waves I & II, 1994–1996; Wave III, 2001–2002; Wave IV, 2007–2009; Wave V, 2016–2018 [machine-readable data file and documentation], 2018.
- Aljoscha Janssen and Xuan Zhang. Retail Pharmacies and Drug Diversion during the Opioid Epidemic. *American Economic Review*, 113(1):1–33, 2023. doi:[10.1257/aer.20210357](https://doi.org/10.1257/aer.20210357).
- Juanna Schrøter Joensen and Helena Skyt Nielsen. Spillovers in Education Choice. *Journal of Public Economics*, 157:158–183, 2018. doi:[10.1016/j.jpubeco.2017.10.006](https://doi.org/10.1016/j.jpubeco.2017.10.006).

- Nazleen F Khan, Brian T Bateman, Joan E Landon, and Joshua J Gagne. Association of Opioid Overdose with Opioid Prescriptions to Family Members. *JAMA internal medicine*, 179(9):1186–1192, 2019. doi:[10.1001/jamainternmed.2019.1064](https://doi.org/10.1001/jamainternmed.2019.1064).
- Seongeun Kim, Michele Tertilt, and Minchul Yum. Status Externalities in Education and Low Birth Rates in Korea. *American Economic Review*, 114(6):1576–1611, 2024. doi:[10.1257/aer.20220583](https://doi.org/10.1257/aer.20220583).
- Michael Kremer and Dan Levy. Peer Effects and Alcohol Use among College Students. *Journal of Economic Perspectives*, 22(3):189–206, 2008. doi:[10.1257/jep.22.3.189](https://doi.org/10.1257/jep.22.3.189).
- Byungkyu Lee, Wanying Zhao, Kai-Cheng Yang, Yong-Yeol Ahn, and Brea L Perry. Systematic Evaluation of State Policy Interventions Targeting the US Opioid Epidemic, 2007-2018. *JAMA network open*, 4(2):e2036687–e2036687, 2021. doi:[10.1001/jamanetworkopen.2020.36687](https://doi.org/10.1001/jamanetworkopen.2020.36687).
- Petter Lundborg. Having the Wrong Friends? Peer Effects in Adolescent Substance Use. *Journal of Health Economics*, 25(2):214–233, 2006. doi:[10.1016/j.jhealeco.2005.02.001](https://doi.org/10.1016/j.jhealeco.2005.02.001).
- Kai Mäcke and Stefan Ruenzi. Friends with Drugs: The Role of Social Networks in the Opioid Epidemic. *Available at SSRN 4104796*, 2022. doi:[10.2139/ssrn.4104796](https://doi.org/10.2139/ssrn.4104796).
- Johanna Catherine Maclean, Justine Mallatt, Christopher J Ruhm, and Kosali Simon. The Opioid Crisis, Health, Healthcare, and Crime: A Review of Quasi-Experimental Economic Studies. *The Annals of the American Academy of Political and Social Science*, 703(1):15–49, 2022. doi:[10.1177/00027162221149285](https://doi.org/10.1177/00027162221149285).
- Charles F Manski. Identification of Endogenous Social Effects: The Reflection Problem. *The Review of Economic Studies*, 60(3):531–542, 1993. doi:[10.2307/2298123](https://doi.org/10.2307/2298123).
- Charles F Manski. Economic Analysis of Social Interactions. *Journal of Economic Perspectives*, 14(3):115–136, 2000. doi:[10.1257/jep.14.3.115](https://doi.org/10.1257/jep.14.3.115).
- John Muench, Katie Fankhauser, Robert W Voss, Nathalie Huguet, Daniel M Hartung, Jean O’Malley, Steffani R Bailey, Stuart Cowburn, Dagan Wright, Gordon Barker, et al. Assessment of Opioid Prescribing Patterns in a Large Network of US Community Health Centers, 2009 to 2018. *JAMA network open*, 3(9):e2013431–e2013431, 2020. doi:[10.1001/jamanetworkopen.2020.13431](https://doi.org/10.1001/jamanetworkopen.2020.13431).

- Eleonora Patacchini, Edoardo Rainone, and Yves Zenou. Heterogeneous Peer Effects in Education. *Journal of Economic Behavior & Organization*, 134:190–227, 2017. doi:[10.1016/j.jebo.2016.10.020](https://doi.org/10.1016/j.jebo.2016.10.020).
- Christiern Rose, Jenny Williams, and Anne Line Bretteville-Jensen. Do Peers Support or Subvert Recovery from Substance Use Disorders. Technical report, HEDG, c/o Department of Economics, University of York, 2024.
- Christopher J Ruhm. Drivers of the Fatal Drug Epidemic. *Journal of Health Economics*, 64:25–42, 2019. ISSN 0167-6296. doi:[10.1016/j.jhealeco.2019.01.001](https://doi.org/10.1016/j.jhealeco.2019.01.001).
- Anil Rupasingha, Stephan J Goetz, and David Freshwater. The Production of Social Capital in US Counties. *The journal of socio-economics*, 35(1):83–101, 2006. doi:[10.1016/j.socec.2005.11.001](https://doi.org/10.1016/j.socec.2005.11.001).
- Marissa J Seamans, Timothy S Carey, Daniel J Westreich, Stephen R Cole, Stephanie B Wheeler, G Caleb Alexander, Virginia Pate, and M Alan Brookhart. Association of Household Opioid Availability and Prescription Opioid Initiation among Household Members. *JAMA Internal Medicine*, 178(1):102–109, 2018. doi:[10.1001/jamainternmed.2017.7280](https://doi.org/10.1001/jamainternmed.2017.7280).
- Peter Rønø Thingholm. Provider Spill-overs in Opioid Prescription Leniency and Patient Labor Market Outcomes. Economics Working Paper, Department of Economics and Business Economics, Aarhus University, 2023. URL [https://repec.econ.au.dk/repec/afn/wp/23/wp23\\_05.pdf](https://repec.econ.au.dk/repec/afn/wp/23/wp23_05.pdf).
- Michael J Zoorob and Jason L Salemi. Bowling Alone, Dying Together: The Role of Social Capital in Mitigating the Drug Overdose Epidemic in the United States. *Drug and alcohol dependence*, 173:1–9, 2017. doi:[10.1016/j.drugalcdep.2016.12.011](https://doi.org/10.1016/j.drugalcdep.2016.12.011).

# Online Appendix

## Figures and Tables

1





Figure A1: Trajectory of the opioid epidemic

**Source:** Calculations based on data from the National Center on Health Statistics, CDC WONDER.






Figure A3: Examples of networks with low centrality (left) and high centrality (right)




Figure A4: Examples of networks with low density (left) and high density (right)

Table A1: Peer effects on opioid misuse-2SLS robustness

|  | Dep. var.: Prob(Opioid misuse) |                     |                     |                     |                               |                      |                           |
|--|--------------------------------|---------------------|---------------------|---------------------|-------------------------------|----------------------|---------------------------|
|  | (1)                            | (2)                 | (3)                 | (4)                 | (5)                           | (6)                  | (7)                       |
| Any best friend opioid misuse                          | 0.416**<br>(0.206)             | 0.380*<br>(0.203)   | 0.466**<br>(0.195)  | 0.490**<br>(0.211)  | 0.459**<br>(0.189)            | 0.533**<br>(0.222)   | 0.441*<br>(0.238)         |
| Severely injured                                       | 0.122***<br>(0.032)            | 0.121***<br>(0.032) | 0.105***<br>(0.029) | 0.109***<br>(0.030) | 0.097***<br>(0.030)           | 0.115***<br>(0.030)  | 0.103***<br>(0.032)       |
| College  | -0.059**<br>(0.023)            | -0.053**<br>(0.024) | -0.030<br>(0.023)   | -0.043*<br>(0.025)  | -0.033<br>(0.026)             | -0.033<br>(0.026)    | -0.028<br>(0.031)         |
| Female   | -0.029<br>(0.019)              | -0.036*<br>(0.018)  | -0.039*<br>(0.020)  | -0.038*<br>(0.019)  | -0.014<br>(0.023)             | -0.029<br>(0.027)    | -0.041*<br>(0.022)        |
| Age  | -0.012<br>(0.009)              | -0.016*<br>(0.009)  | -0.009<br>(0.007)   | -0.010<br>(0.007)   | -0.006<br>(0.007)             | -0.007<br>(0.007)    | -0.013<br>(0.008)         |
| Hispanic   | 0.017<br>(0.047)               | 0.017<br>(0.048)    | 0.003<br>(0.040)    | 0.003<br>(0.049)    | 0.002<br>(0.040)              | 0.010<br>(0.040)     | -0.030<br>(0.055)         |
| African American                                       | -0.030<br>(0.046)              | -0.025<br>(0.046)   | 0.004<br>(0.036)    | 0.009<br>(0.080)    | 0.006<br>(0.035)              | 0.015<br>(0.039)     | 0.025<br>(0.050)          |
| Ever diagn. depressed                                  | 0.090*<br>(0.046)              | 0.082*<br>(0.043)   | 0.070*<br>(0.041)   | 0.067<br>(0.041)    | 0.056<br>(0.041)              | 0.072*<br>(0.042)    | 0.054<br>(0.048)          |
| Ever diagn. post-traumatic stress                      | 0.060<br>(0.065)               | 0.074<br>(0.066)    | 0.103<br>(0.064)    | 0.108*<br>(0.064)   | 0.089<br>(0.063)              | 0.129*<br>(0.069)    | 0.039<br>(0.073)          |
| Ever diagn. anxiety                                    | 0.021<br>(0.047)               | 0.035<br>(0.043)    | 0.036<br>(0.043)    | 0.041<br>(0.044)    | 0.031<br>(0.043)              | 0.037<br>(0.046)     | 0.053<br>(0.047)          |
| Cigarettes avail. at parental home in WI               | 0.020<br>(0.030)               | 0.023<br>(0.031)    | 0.013<br>(0.031)    | 0.016<br>(0.033)    | 0.018<br>(0.032)              | 0.011<br>(0.029)     | 0.049<br>(0.035)          |
| Alcohol avail. at parental home in WI                  | 0.030<br>(0.029)               | 0.028<br>(0.027)    | 0.019<br>(0.028)    | 0.017<br>(0.028)    | 0.018<br>(0.028)              | 0.021<br>(0.032)     | 0.008<br>(0.040)          |
| Drugs avail. at parental home in WI                    |                                |                     | 0.142*<br>(0.076)   | 0.145*<br>(0.077)   |                               |                      |                           |
| % college educated best friends                        |                                |                     |                     | 0.036<br>(0.031)    |                               |                      |                           |
| % Hispanic best friends                                |                                |                     |                     | 0.020<br>(0.051)    |                               |                      |                           |
| % African American best friends                        |                                |                     |                     | 0.008<br>(0.077)    |                               |                      |                           |
| % best friends with cigarettes avail. at parental home |                                |                     |                     | -0.014<br>(0.029)   |                               |                      |                           |
| % best friends with alcohol avail. at parental home    |                                |                     |                     | 0.023<br>(0.029)    |                               |                      |                           |
| % best friends with drugs avail. at parental home      |                                |                     |                     | -0.031<br>(0.070)   |                               |                      |                           |
| Extraversion   |                                |                     |                     |                     | -0.001<br>(0.004)             |                      |                           |
| Neuroticism  |                                |                     |                     |                     | 0.005<br>(0.004)              |                      |                           |
| Agreeableness  |                                |                     |                     |                     | 0.003<br>(0.005)              |                      |                           |
| Conscientiousness                                      |                                |                     |                     |                     | -0.019***<br>(0.005)          |                      |                           |
| Openness   |                                |                     |                     |                     | 0.014***<br>(0.005)           |                      |                           |
| Risk aversion  |                                |                     |                     |                     | -0.023**<br>(0.011)           |                      |                           |
| Maternal education in WI                               |                                |                     |                     |                     |                               |                      | 0.008<br>(0.021)          |
| Gross Hhd income in thousand \$ in WI                  |                                |                     |                     |                     |                               |                      | 0.001**<br>(0.000)        |
| Live with both parents in WI                           |                                |                     |                     |                     |                               |                      | -0.053*<br>(0.028)        |
| Observations   | 2,826                          | 2,823               | 2,824               | 2,804               | 2,807                         | 2,767                | 2,151                     |
| FE   | School                         | School and State    | State               | State               | State                         | State                | State                     |
| Description  | Different FE                   | Different FE        | Drug avail. control | Peer char. controls | Big5 & risk aversion controls | Occupational dummies | Family of origin controls |
| Kleibergen-Paap Wald rk F statistic                    | 16.37                          | 17.39               | 18.47               | 16.47               | 18.97                         | 16.72                | 15.12                     |

**Note:** The estimated coefficients of equation 2 and the F-statistic of equation 3 with different sets of fixed effects, control variables, and sample restrictions (columns 1-7). Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \*p<0.10; \*\*p<0.05; \*\*\*p<0.01.



Table A2: Peer effects on opioid misuse-2SLS different measures of peer behavior

|                                     | Dep. var.: Opioid misuse       |                               |                           |                          |
|-------------------------------------|--------------------------------|-------------------------------|---------------------------|--------------------------|
|                                     | (1)                            | (2)                           | (3)                       | (4)                      |
| Best friend opioid misuse           | 0.472**<br>(0.196)             | 0.544**<br>(0.266)            | 0.451**<br>(0.193)        | 0.530**<br>(0.235)       |
| Observations                        | 2,826                          | 2,826                         | 2,826                     | 3,257                    |
| Controls                            | Yes                            | Yes                           | Yes                       | Yes                      |
| FE                                  | State                          | State                         | State                     | State                    |
| Description                         | Any best friend<br>(Benchmark) | Percentage of<br>best friends | Number of<br>best friends | Dyads of<br>best friends |
| Kleibergen-Paap Wald rk F statistic | 18.44                          | 12.72                         | 16.98                     | 12.10                    |

**Note:** The estimated coefficients of equation 2 and the F-statistic of equation 3 with different measures of peer behavior. Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \*p<0.10; \*\*p<0.05; \*\*\*p<0.01. See Table 4, column 6 for the list of controls.

Table A3: Share of opioid misusers by education and race

| By education |             | By race            |        |
|--------------|-------------|--------------------|--------|
| College      | Non-College | Non-Hispanic white | Others |
| (1)          | (2)         | (3)                | (4)    |
| 0.147        | 0.182       | 0.193              | 0.063  |

**Note:** Characteristics of individuals in the Add Health regression sample. Survey weights are used.

Table A4: Peer effects on opioid misuse-2SLS by type of friends

|                                     | Dep. var.: Prob(Opioid misuse) |                   |                               |                            |                        |                             |                                  |                              |                                   |
|-------------------------------------|--------------------------------|-------------------|-------------------------------|----------------------------|------------------------|-----------------------------|----------------------------------|------------------------------|-----------------------------------|
|                                     | Best friends                   | Other friends     | Best friends<br>in WI and WII | Best friends<br>only in WI | Same gender<br>friends | Same county<br>best friends | Different county<br>best friends | Same county<br>other friends | Different county<br>other friends |
|                                     | (1)                            | (2)               | (3)                           | (4)                        | (5)                    | (6)                         | (7)                              | (8)                          | (9)                               |
| Any best friend opioid misuse       | 0.472**<br>(0.196)             | -0.196<br>(0.155) | 0.605**<br>(0.292)            | -0.233<br>(0.179)          | 0.746**<br>(0.349)     | 0.916**<br>(0.378)          | -0.209<br>(0.390)                | -0.501*<br>(0.275)           | -0.255<br>(0.317)                 |
| Observations                        | 2,826                          | 1,873             | 935                           | 1,034                      | 2,127                  | 1,330                       | 1,304                            | 1,130                        | 1,033                             |
| State FE                            | Yes                            | Yes               | Yes                           | Yes                        | Yes                    | Yes                         | Yes                              | Yes                          | Yes                               |
| Mean                                | 0.169                          | 0.190             | 0.181                         | 0.162                      | 0.164                  | 0.160                       | 0.169                            | 0.167                        | 0.203                             |
| Kleibergen-Paap Wald rk F statistic | 18.44                          | 35.76             | 12.27                         | 13.19                      | 10.97                  | 8.407                       | 10.13                            | 11.94                        | 11.55                             |

**Note:** Estimated coefficients of equation 2 and F-statistic of equation 3 for best (first nominated) friends (column 1); other (second, third, fourth or fifth nominated) friends (column 2); for best friends nominated both in Wave I and Wave II (column 3); for best friends nominated only in Wave I (column 4); for best friends of the same gender as the respondent (column 5); for best friends residing in the same or different county as the respondent (columns 6 and 7); and for other friends residing in the same or different county as the respondent (columns 8 and 9). Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \*p<0.10; \*\*p<0.05; \*\*\*p<0.01. See Table 4, column 6 for the list of controls.

Table A5: Peer effects on opioid misuse-2SLS by network characteristics

|                                     | Own Bonacich Centrality |                    | Best friends' Bonacich Centrality |                     | Networks' reach  |                    | Networks' relative density |                    |
|-------------------------------------|-------------------------|--------------------|-----------------------------------|---------------------|------------------|--------------------|----------------------------|--------------------|
|                                     | Low<br>(1)              | High<br>(2)        | Low<br>(3)                        | High<br>(4)         | Low<br>(5)       | High<br>(6)        | Low<br>(7)                 | High<br>(8)        |
| Any best friend opioid misuse       | 0.177<br>(0.302)        | 0.659**<br>(0.297) | 0.379<br>(0.379)                  | 0.525***<br>(0.195) | 0.332<br>(0.257) | 0.665**<br>(0.321) | 0.362<br>(0.286)           | 0.535**<br>(0.241) |
| Observations                        | 1,078                   | 1,743              | 1,141                             | 1,677               | 1,103            | 1,715              | 1,406                      | 1,414              |
| State FE                            | Yes                     | Yes                | Yes                               | Yes                 | Yes              | Yes                | Yes                        | Yes                |
| Mean                                | 0.161                   | 0.173              | 0.157                             | 0.176               | 0.161            | 0.175              | 0.177                      | 0.164              |
| Kleibergen-Paap Wald rk F statistic | 9.667                   | 11.16              | 4.794                             | 17.75               | 8.917            | 13.10              | 8.270                      | 11.49              |

**Note:** Estimates for individuals that are less or more central in the entire friendship network (own Bonacich centrality below or above the mean, columns 1 and 2); for individuals whose best friends are less or more central in the entire friendship network (best friends' Bonacich centrality below or above the mean, columns 3 and 4); for individuals with a low or high reach (maximum number of individuals a node can connect within the entire friendship network is below or above the mean, columns 5 and 6); and for individuals in relatively less or more dense friendship networks (columns 7 and 8). Relative density is measured as the number of actual ties in the total friendship network divided by the maximum possible density given an out-degree of 10 (each respondent could nominate at most 10 friends). Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \*p<0.10; \*\*p<0.05; \*\*\*p<0.01. See Table 4, column 6 for the list of controls.

Table A6: Peer effects on opioid misuse-2SLS by local level conditions

| Dep. var.: Prob(Opioid misuse)      |                      |                    |                        |                   |   |                    |                      |                  |  |                    |  |                   |
|-------------------------------------|----------------------|--------------------|------------------------|-------------------|---|--------------------|----------------------|------------------|--|--------------------|--|-------------------|
|                                     | Opioid dispense rate |                    | Naloxone accessibility |                   | Average number of poor mental health days per month |                    | Social capital index |                  | Number of mental health facilities within 20 miles |                    | Number of physicians per 100k population |                   |
|                                     | Low (1)              | High (2)           | Low (3)                | High (4)          | Low (5)   | High (6)           | Low (7)              | High (8)         | Low (9)  | High (10)          | Low (11)                                 | High (12)         |
| Any best friend opioid misuse       | 0.108<br>(0.192)     | 0.625**<br>(0.263) | 0.187<br>(0.240)       | 0.673*<br>(0.378) | 0.320<br>(0.210)                                    | 0.628**<br>(0.262) | 0.585**<br>(0.287)   | 0.237<br>(0.251) | 0.134<br>(0.192)                                   | 0.946**<br>(0.443) | 0.191<br>(0.224)                         | 0.779*<br>(0.462) |
| Observations                        | 979                  | 948                | 1,153                  | 770               | 1,016   | 911                | 905                  | 1,019            | 966  | 960                | 1,007                                    | 912               |
| Controls                            | Yes                  | Yes                | Yes                    | Yes               | Yes   | Yes                | Yes                  | Yes              | Yes  | Yes                | Yes                                      | Yes               |
| State FE                            | Yes                  | Yes                | Yes                    | Yes               | Yes   | Yes                | Yes                  | Yes              | Yes  | Yes                | Yes                                      | Yes               |
| Kleibergen-Paap Wald rk F statistic | 21.16                | 9.701              | 17.38                  | 9.527             | 18.72   | 9.502              | 10.95                | 11.92            | 12.57  | 6.415              | 19.44                                    | 5.264             |

**Note:** Estimates for individuals who reside in states with opioid dispense rates below or above the median (columns 1 and 2); for individuals who reside in states that implemented naloxone laws earlier or later (columns 3 and 4); for individuals who reside in counties with an average number of poor mental health days per month below or above the median (columns 5 and 6); for individuals who reside in counties with a social capital index in 2014 below or above the median (columns 7 and 8); for individuals who reside in locations with less or more than 13 mental health facilities within 20 miles (columns 9 and 10); for individuals who reside in counties with less or more than 70 primary-care physicians per 100,000 population (columns 11 and 12). The location of residence is as of the date of the Wave V interview (that took place in the period 2016-2018). Robust standard errors in parentheses are clustered at the school level. Survey weights are used. \*p<0.10; \*\*p<0.05; \*\*\*p<0.01. See Table 4, column 6 for the list of controls.